arXiv:1209.4073v2 [math.DS] 24 Sep 2012

SECOND ORDER ERGODIC THEOREM FOR SELF-SIMILAR TILING SYSTEMS

KONSTANTIN MEDYNETS AND BORIS SOLOMYAK

Abstract. We consider infinite measure-preserving non-primitive self-similar tiling systems in Euclidean space $\mathbb{R}^d$. We establish the second-order ergodic theorem for such systems. The exponent of the averaging sequence is the Hausdorff dimension of the graph-directed set associated with the substitution rule.

1. Introduction

Let $\mathbb{X} = (X, \mu, \mathbb{T})$ be a conservative, ergodic, measure preserving dynamical system of a σ-finite measure space. If $f \in L^1(X, \mu)$, then Hopf's ratio ergodic theorem says that the growth of $S_n f(x) = f(x) + \cdots + f(T^{n-1}x)$ is independent of f in the sense that if $g \in L^1(X, \mu)$ with $\int g\, d\mu \neq 0$, then

$$\frac{S_n f(x)}{S_n g(x)} \to \frac{\int f\, d\mu}{\int g\, d\mu} \text{ for } \mu\text{-a.e. } x \in X.$$

It turns out that due to the measure μ being infinite, it is impossible to replace functions $S_n g(x)$ by constants $\{a_n\}$ [A, Theorem 2.4.2]. However, it was observed earlier [Fi1, ADF, LS] that for some systems, the ratios $S_n f(x)/a_n$ (for some choice of a_n) still converge to $\int f\, d\mu$ though in a weaker sense (second-order averages). The asymptotic behavior of the sequence $\{a_n\}$ is an invariant of the dynamical system.

The main result of the present paper is the following ergodic theorem showing that for self-similar tilings the sequence $\{a_n\}$ can be chosen as $\{n^{\alpha+1}\}$ where α is an intrinsic parameter of the system reflecting self-similarity (the precise statements, with all technical assumptions, are Theorem 5.4 and Theorem 6.11).

Theorem 1.1. *(i) Let $\mathbb{X} = (\Omega, \mu, \mathbb{R}^d)$ be a non-primitive self-similar substitution tiling system preserving an infinite ergodic measure μ. Assume that the measure μ is non-zero and finite on some open subset of Ω. Then there exist positive parameters α and c such that for μ-almost every tiling $\mathcal{T} \in \Omega$ and for every function $f \in L^1(\Omega, \mu)$, we have*

$$\lim_{t\to\infty} \frac{1}{\log(t)} \int_0^t \frac{\int_{B_R} f(\mathcal{T} - u)\, du}{cR^\alpha} \frac{dR}{R} = \int_\Omega f d\mu. \tag{1.1}$$

Date: September 25, 2012.

B. S. is supported in part by NSF grant DMS-0968879.

Here B_R is the ball of radius R centered at the origin.

(ii) The parameters α and c are invariants of the measure-theoretic isomorphism of the system $\mathbb{X}$.

(iii) This result is also valid for a large class of one-dimensional symbolic substitution systems with integrals in the left-hand side replaced by the corresponding sums.

Our work was originally inspired by A. Fisher's paper [Fi1], where he obtained a similar ergodic theorem for a single substitution system (X_σ, S) generated by the map $\sigma(0) = 000$ and $\sigma(1) = 101$. Iterating the map $\sigma^n(1)$, $n \geq 1$, we get a sequence where the appearances of 1's and 0's resemble the process of constructing the middle-thirds Cantor set. Fisher used this analogy to show that for any function $f \in L^1(X_\sigma, \mu)$ and μ-a.e point $x \in X_\sigma$,

$$\lim_{n\to\infty} \frac{1}{\log(n)} \sum_{k=1}^{n} \frac{\sum_{i=0}^{k-1} f(S^i x)}{ck^\alpha} \cdot \frac{1}{k} = \int_{X_\sigma} f d\mu,$$

where μ is the unique S-invariant measure μ on X_σ with $\mu([1]) = 1$. Here $\alpha = \log(2)/\log(3)$ is the Hausdorff dimension and c is the average density of the middle-thirds Cantor set. The middle-thirds Cantor set arising in the study of the substitution σ is a special case of the graph-directed set that one can associate to every self-similar tiling system (Section 3.1). In the proof of Theorem 1.1, we show the parameters α and c always arise as the Hausdorff dimension and the average density, respectively, of the associated graph-directed sets.

It is well-known that primitive substitution dynamical systems (both symbolic and tiling versions) are uniquely ergodic. Non-primitive substitutions and their invariant measures have been recently studied in [Y, BKM, BKMS1, HY], where it was shown, in particular, that for a large class of non-primitive symbolic substitutions infinite (σ-finite) invariant measures appear naturally. This has also been extended to the tiling setting in [CS]. The present work continues this line of research, focusing on ergodic properties of the resulting infinite measure-preserving transformations.

We observe that the second-order ergodic theorem is not a universal result. Its validity depends on intrinsic properties of the dynamical system in question. We refer the reader to the paper [ADF] of Aaronson-Denker-Fisher for the discussion of second order ergodic theorems for Markov shifts. We also mention the paper [LS] of Ledrappier and Sarig establishing the second order ergodic theorem for certain horocycle flows.

The structure of the paper is the following. In Section 2 we give the definition of the substitution tiling system $(\Omega, \mathbb{R}^d)$ generated by a tile substitution $\mathcal{G}$. We then recall the classification of infinite ergodic invariant measures established in [CS]. In subsection 2.3 we explicitly state technical assumptions on the tiling system needed for Theorem 1.1.

In Section 3 we show how to associate a graph-directed set to the substitution rule $\mathcal{G}$. In the following sections we use this fractal to count frequencies of prototiles in tilings of Ω.

In Section 4, we show that the transverse dynamical system (generated by iterations of the substitution rule) is measure-theoretically isomorphic to a Markov chain.

Sections 5 and 6 are devoted to the proof of Theorem 1.1. As a corollary, we establish that almost every sequence from a substitution space admits a non-zero α-dimensional density, where the parameter α comes from Theorem 1.1. Section 7 contains a few examples and open questions.

2. Tiling Dynamical Systems

In this section we fix our notation and present necessary definitions from the theory of tiling dynamical systems. We mostly follow conventions of [CS].

2.1. **Tiling Space.**

Fix a finite alphabet $\mathcal{A}$ and an integer $d \geq 1$. By a *tile* in $\mathbb{R}^d$ we mean a pair $T = (F, i)$ of a compact set F that is the closure of its interior, and a letter (label) $i \in \mathcal{A}$. Two geometrically identical sets labeled by different letters are treated as distinct tiles. The set F will be called the *support* of the tile T, in symbols, $\mathrm{supp}(T) = F$. A *tiling* is a family of tiles $\mathcal{T}$ such that $\mathbb{R}^d = \cup\{\mathrm{supp}(T) : T \in \mathcal{T}\}$ and distinct tiles have disjoint interiors. A *patch* is a finite set of tiles with disjoint interiors. The *support of a patch* P is the set $\mathrm{supp}(P) = \cup\{\mathrm{supp}(T) : T \in P\}$. If $\mathcal{T}$ is a tiling, its finite subsets are called $\mathcal{T}$*-patches.*

The translate of a tile $T = (F, i)$ by a vector $u \in \mathbb{R}^d$ is the tile $T + u = (F + u, i)$. Similarly, a translate of a patch P by $u \in \mathbb{R}^d$ is the patch $P + u = \{T + u : T \in P\}$. We say that two patches P_1 and P_2 are *translationally equivalent* if $P_1 = P_2 + u$ for some vector $u \in \mathbb{R}^d$.

Definition 2.1. Let $\mathcal{A}$ be a finite set of tiles in $\mathbb{R}^d$ such that distinct tiles from $\mathcal{A}$ are not translationally equivalent. Tiles from the set $\mathcal{A}$ are called *prototiles.* We will call translations of prototiles $\mathcal{A}$*-tiles.*

Denote by $\mathcal{A}^+$ the set of patches made of translates of tiles from $\mathcal{A}$. We assume that every prototile $T \in \mathcal{A}$ is centered at the origin, in the sense that $\mathbf{0} \in \mathrm{int}(\mathrm{supp}(T))$.

Let φ be an expanding linear transformation in $\mathbb{R}^d$. A map $\mathcal{G} : \mathcal{A} \to \mathcal{A}^+$ is called a *tile substitution with expansion* φ if

$$\mathrm{supp}(\mathcal{G}(T)) = \varphi(\mathrm{supp}(T)) \text{ for every tile } T \in \mathcal{A}. \tag{2.1}$$

In other words, the substitution $\mathcal{G}$ shows how to subdivide the inflated tile $\varphi(\mathrm{supp}(T))$ into translates of prototiles. The tile substitution can be written explicitly as follows:

$$\mathcal{G}(T) = \bigcup_{T' \in \mathcal{A}} \{T' + u : \ u \in \mathcal{D}_{T,T'}\} \ \text{ for all } \ T \in \mathcal{A}, \tag{2.2}$$

where $\mathcal{D}_{T,T'}$ is a finite (possibly empty) subset of $\mathbb{R}^d$, the tiles in the right-hand side have disjoint interiors, and

$$\varphi(\mathrm{supp}(T)) = \bigcup_{T'\in\mathcal{A}} \bigcup_{u\in\mathcal{D}_{T,T'}} (\mathrm{supp}(T') + u). \tag{2.3}$$

The substitution $\mathcal{G}$ is extended to translates of prototiles by $\mathcal{G}(T+u) = \mathcal{G}(T) + \varphi(u)$; and to patches by $\mathcal{G}(P) = \cup\{\mathcal{G}(T) : T \in P\}$. The linearity of φ and the equation (2.1) imply that the patch $\mathcal{G}(P)$ is well-defined.

Remark 2.2. In this paper we restrict ourselves to the *self-similar* case, i.e. $\varphi = \lambda \cdot O$, where O is an orthogonal matrix and $\lambda > 1$. We refer to the corresponding $\mathcal{G}$ as *self-similar tiling substitution.* The more general case of an arbitrary expansion map φ, referred to as *self-affine*, is not covered by our main results.

Definition 2.3. For a given tiling substitution $\mathcal{G} : \mathcal{A} \to \mathcal{A}^+$, let $M = (m_{A,B})_{A,B\in\mathcal{A}}$ be the matrix with $m_{A,B}$ being the number of translates of the prototile A in the patch $\mathcal{G}(B)$ (i.e. $m_{A,B} = \#\mathcal{D}_{A,B}$). The matrix M is called the *substitution matrix of* $\mathcal{G}$.

The substitution is called *primitive* if some power of the substitution matrix has only positive entries. We emphasize that our focus is on the non-primitive case.

Definition 2.4. Let $\mathcal{G} : \mathcal{A} \to \mathcal{A}^+$ be a tile substitution. Denote by $\Omega_{\mathcal{G}}$ the set of all tilings of $\mathbb{R}^d$ by tiles from $\mathcal{A}$ such that $\mathcal{T} \in \Omega_{\mathcal{G}}$ if every $\mathcal{T}$-patch is a subpatch of $\mathcal{G}^n(T) + u$ for some $T \in \mathcal{A}$, $u \in \mathbb{R}^d$, and $n \geq 1$. The set $\Omega_{\mathcal{G}}$ is called the *tiling space* corresponding to the substitution $\mathcal{G}$.

Let $\|\cdot\|$ be the Euclidean norm on $\mathbb{R}^d$. For $x \in \mathbb{R}^d$ and $R > 0$, set $B_R(x) = \{u \in \mathbb{R}^d : \|u - x\| \leq R\}$. We will write B_R for $B_R(\mathbf{0})$. For a compact set K and a tiling $\mathcal{T}$, denote by $\mathcal{T}[[K]]$ the set of all $\mathcal{T}$-patches P with $K \subset \mathrm{supp}(P)$. Define a metric ρ on the space $\Omega_{\mathcal{G}}$ as follows. Given tilings $\mathcal{T}', \mathcal{T}'' \in \Omega_{\mathcal{G}}$, let $\rho(\mathcal{T}', \mathcal{T}'')$ be the minimum of $2^{-1/2}$ and

$$\inf\{r > 0 : \exists g \in B_r, P' \in \mathcal{T}'[[B_{1/r}]], P'' \in \mathcal{T}''[[B_{1/r}]] \text{ such that } P' - g = P''\}.$$

With respect to the topology generated by ρ, two tilings are close to each other if they agree on a large ball around the origin after a small translation. The cut-off parameter $2^{-1/2}$ is needed to fulfill the triangle inequality.

Definition 2.5. (1) We say that the tiling system $\Omega_{\mathcal{G}}$ satisfies the *finite pattern condition* (FPC) if for every tiling $\mathcal{T} \in \Omega_{\mathcal{G}}$ and $R > 0$, there are only finitely many $\mathcal{T}$-patches of diameter less than R up to translation equivalence. (Note that *finite local complexity* and *translational finiteness* are used synonymously with FPC).

(2) The tiling substitution $\mathcal{G}$ is called *admissible* if for every prototile $T \in \mathcal{A}$ there exists a tile $\mathcal{T} \in \Omega_{\mathcal{G}}$ such that $T \in \mathcal{T}$.

The FPC assumption implies the following result. The proof can be found, for example, in [RW, Lemma 2].

Proposition 2.6. *If the tiling system has the FPC property, then the set* $\Omega_{\mathcal{G}}$ *is compact with respect to the topology generated by the metric* ρ. *The action of the group* $\mathbb{R}^d$ *by translations on* $\Omega_{\mathcal{G}}$ *is continuous.*

The following result shows that any tiling in $\Omega_{\mathcal{G}}$ has a preimage under the map $\mathcal{G}$, see [CS, Lemma 2.8].

Proposition 2.7. *If* $\mathcal{G}$ *is admissible, then the map* $\mathcal{G} : \Omega_{\mathcal{G}} \to \Omega_{\mathcal{G}}$ *is a continuous surjection.*

One of the important issues in the theory of substitutions is to understand when the map $\mathcal{G}$ is invertible. This property is sometimes referred to as *recognizability.* It is equivalent to non-periodicity of the tiling space for primitive tile substitutions [So2], but the extension to the non-primitive case is by no means trivial. To state conditions ensuring the invertibility of the map $\mathcal{G}$, we need the following definitions.

Definition 2.8. (1) A tiling $\mathcal{T} \in \Omega_{\mathcal{G}}$ is *periodic* if there is a non-zero vector $u \in \mathbb{R}^d$ such that $\mathcal{T} = \mathcal{T} + u$. A tile substitution $\mathcal{G}$ is called *non-periodic* if the set $\Omega_{\mathcal{G}}$ has no periodic tilings.

(2) A *minimal component* of the system $(\Omega_{\mathcal{G}}, \mathbb{R}^d)$ is an $\mathbb{R}^d$-invariant set that contains no proper closed invariant subsets.

(3) Denote by $\mathcal{A}_{\text{per}}$ the set of all tiles $T \in \mathcal{A}$ that occur in tilings from periodic minimal components. Set $\mathcal{A}_{\text{nonp}} = \mathcal{A} \setminus \mathcal{A}_{\text{per}}$. For a patch P, let $P|_{\text{nonp}}$ be the subpatch of all $\mathcal{A}_{\text{nonp}}$-tiles in P.

(4) A substitution $\mathcal{G}$ is said to *satisfy the non-periodic border condition* if for every tile $T \in \mathcal{A}_{\text{nonp}}$, the $\mathbb{R}^d$-boundary of the support of the patch $\mathcal{G}(T)$ is contained in the set $\text{supp}(\mathcal{G}(T)|_{\text{nonp}})$.

For the proof of the following result see [CS, Theorem 4.1 and 4.4].

Theorem 2.9. *(1) If the dynamical system* $(\Omega_{\mathcal{G}}, \mathbb{R}^d)$ *has no periodic orbits, then the substitution* $\mathcal{G} : \Omega_{\mathcal{G}} \to \Omega_{\mathcal{G}}$ *is a homeomorphism.*

(2) Assume that a substitution $\mathcal{G}$ *satisfies the non-periodic border condition. Then for every tiling* $\mathcal{T} \in \Omega_{\mathcal{G}}$ *that contains a tile from* $\mathcal{A}_{nonp}$ *there exists a unique tiling* $\mathcal{T}'$ *such that* $\mathcal{G}(\mathcal{T}') = \mathcal{T}$.

Reordering the letters in the alphabet $\mathcal{A}$ and replacing the substitution $\mathcal{G}$ with its higher powers $\mathcal{G}^k$ if needed, the substitution matrix can be transformed into the following form:

$$M(\mathcal{G}) = \begin{pmatrix} F_1 & 0 & \cdots & 0 & X_{1,s+1} & \cdots & X_{1,m} \\ 0 & F_2 & \cdots & 0 & X_{2,s+1} & \cdots & X_{2,m} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & \cdots & F_s & X_{s,s+1} & \cdots & X_{s,m} \\ 0 & 0 & \cdots & 0 & F_{s+1} & \cdots & X_{s+1,m} \\ \vdots & \vdots & \cdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & 0 & \cdots & F_m \end{pmatrix} \tag{2.4}$$

The square matrices F_i on the main diagonal are either zero matrices or contain only strictly positive entries. For any fixed $j = s+1, ..., m$, at least one of the matrices $X_{k,j}$ is non-zero. The block-triangular form (2.4) allows us to give an effective description of minimal components of the system. We refer the reader to the papers [BKMS1] and [CS] for more details on the structure of minimal components of substitution and tiling systems.

For each $i = 1, \ldots, m$, denote by $\mathcal{A}_i$ the set of prototiles corresponding to the block F_i. Denote by Ω_i the set of tilings $\mathcal{T} \in \Omega_{\mathcal{G}}$ whose patches are subpatches of $\mathcal{G}^n(T)$, $n \geq 0$, with $T \in \mathcal{A}_i$. The sets $\{\Omega_1, \ldots, \Omega_s\}$ are the only minimal components of the tiling system, see [CS, Lemma 2.10] for the proof.

2.2. **Invariant Measures.**

Definition 2.10. (1) A measure μ on $\Omega_{\mathcal{G}}$ is called *invariant* if $\mu(U - u) = \mu(U)$ for every $u \in \mathbb{R}^d$ and every Borel set $U \subset \Omega_{\mathcal{G}}$. An invariant measure μ is called *ergodic* if whenever a Borel set X is translation-invariant, i.e. $X - u = X$ for every $u \in \mathbb{R}^d$, either $\mu(X) = 0$ or $\mu(\Omega_{\mathcal{G}} \setminus X) = 0$.

(2) By the *transversal* of $\Omega_{\mathcal{G}}$ we mean the family of all tilings $\mathcal{T} \in \Omega_{\mathcal{G}}$ such that one of the $\mathcal{T}$-tiles is exactly a prototile from $\mathcal{A}$. Recall that each prototile contains the origin in the interior of its support. Throughout the paper, the transversal will be denoted by $\Gamma \subset \Omega_{\mathcal{G}}$.

(3) A *transverse measure* is a Borel measure ν on Γ such that $\nu(U) = \nu(U - u)$ for every Borel subset $U \subset \Gamma$ and $u \in \mathbb{R}^d$ for which $U - u \subset \Gamma$.

Proposition 2.11. *There is a one-to-one correspondence between finite (resp. σ-finite) transverse measures and finite (resp. σ-finite) invariant measures* [CS, Section 7].

Consider the transversal Γ. For a prototile $T \in \mathcal{A}$, set

$$\Gamma_T = \{\mathcal{T} \in \Gamma : T \in \mathcal{T}\}.$$

Then $\Gamma = \coprod_{T \in \mathcal{A}} \Gamma_T$ is a disjoint union. The following result provides a description of "natural" σ-finite ergodic measures, for the proof see Theorems 3.1 and 5.22 in [CS].

Theorem 2.12. *(i) Each finite ergodic measure is supported by one of the minimal components* $\{\Omega_1, \ldots, \Omega_s\}$.

(ii) Let $i \in \{s+1, \dots, m\}$ *be such that the matrix* F_i *is nonzero and there exist* $A \in \mathcal{A}_i$ *and* $n > 0$ *such that a translate of* A *appears in the interior of* $\mathcal{G}^n(A)$. *Then there exists a unique (up to scaling) invariant ergodic* σ*-finite measure* μ *supported by* Ω_i *such that* $0 < \mu^{\mathrm{tr}}(\Gamma_C) < \infty$ *for some (and, in fact, for all) prototile* $C \in \mathcal{A}_i$, *where* μ^{tr} *is the transverse measure corresponding to* μ. *Moreover, the vector* $(\mu^{\mathrm{tr}}(\Gamma_C))_{C \in \mathcal{A}_i}$ *is a right Perron-Frobenius eigenvector of* F_i.

Remark 2.13. Denote by $\mathcal{L}^d$ the Lebesgue measure on $\mathbb{R}^d$. Observe that the substitution matrix $M(\mathcal{G})$ has a strictly positive left eigenvector $(\mathcal{L}^d(\mathrm{supp}(T)))_{T \in \mathcal{A}}$, corresponding to the eigenvalue $\lambda = \rho(A)$. This follows from (2.1) and the fact that the tile boundaries have zero d-dimensional Lebesgue measure. (The latter is proved e.g. in [P, Prop. 1.2] by B. Praggastis; she does not assume primitivity there.) Note also that $\rho(A_i) = \lambda$ for $i \le s$ and $\rho(A_i) < \lambda$ for $i = s+1, \dots, m$. The latter inequality follows from the existence of strictly positive left eigenvector, see [G, Theorem III.6].

In view of Theorem 2.12, the study of ergodic measures can be reduced to the study of the dynamics on one of the sets Ω_i. This implies that it is enough to consider substitution tiling systems whose incidence matrices have the following form:

$$M(\mathcal{G}) = \begin{pmatrix} A & C \\ 0 & B \end{pmatrix}, \tag{2.5}$$

where A and B are square matrices; B is a primitive matrix; C and A are non-zero matrices. Note that the matrix A *does not have to be primitive* and may contain zero entries.

2.3. **Technical Assumptions.** Here we summarize the assumptions we will be implicitly imposing on the tiling systems in question. Throughout the paper, the symbols $\mathcal{G}$ and $\varphi(x) = \lambda \cdot O(x)$, $\Omega_{\mathcal{G}}$ will be reserved for a self-similar tile substitution, the associated expansion map, and the tiling space, respectively. The set of prototiles corresponding to the matrix B will be denoted by $\mathcal{B}$. Furthermore, we will always assume that the tiling substitution $\mathcal{G} : \mathcal{A} \to \mathcal{A}^+$ satisfies the following conditions:

1. Every prototile is a compact subset of $\mathbb{R}^d$ that is the closure of its interior. Note that this implies that the Hausdorff dimension of the boundary of every prototile is at least $d-1$, see, for example, Corollary IV.2 and Theorem VII.3 in [HuW].
2. The tiling system $\Omega_{\mathcal{G}}$ satisfies the finite pattern condition.
3. The tile substitution $\mathcal{G}$ is admissible.
4. The substitution $\mathcal{G}$ satisfies the non-periodic border condition (see Definition 2.8).
5. The substitution matrix $M(\mathcal{G})$ has the form (2.5), with C non-zero, B primitive, $\rho(B) > 1$.

(6) We have $\alpha := \log(\rho(B))/\log(\lambda) > d-1$. The meaning of α will be clarified in Section 3.1.

Remark 2.14. The admissibility assumption implies that for every prototile $T \in \mathcal{B}$ there is $n > 0$ such that a translate of T occurs in the interior of $\mathcal{G}^n(T)$.

We summarize dynamical properties of tiling systems, which follow from our assumptions.

Proposition 2.15. *The space $\Omega_{\mathcal{G}}$ is compact. The map $\mathcal{G} : \Omega_{\mathcal{G}} \to \Omega_{\mathcal{G}}$ is a continuous surjection that is invertible on non-periodic tilings. The dynamical system $(\Omega_{\mathcal{G}}, \mathbb{R}^d)$ has a unique (up to scaling) infinite σ-finite measure μ such that the corresponding transverse measure is positive and finite on one (equivalently on all) of the sets Γ_T, $T \in \mathcal{B}$.*

2.4. **Hierarchical Structure.** For any tiling $\mathcal{T} \in \Omega_{\mathcal{G}}$, there is a hierarchical sequence of *scaled* tilings $\mathcal{T}^{(k)}$ for $k \in \mathbb{Z}$, such that $\mathcal{T} = \mathcal{T}^{(0)}$ and $\mathcal{T}^{(k-1)}$ is obtained from $\mathcal{T}^{(k)}$ applying the (rescaled) subdivision rule. If $\mathcal{T}$ contains a tile of type $\mathcal{A}_{\rm nonp}$, then we simply have

$$\mathcal{T}^{(k)} = \varphi^k \mathcal{G}^{-k}(\mathcal{T}) \quad \text{for all } k \in \mathbb{Z}. \tag{2.6}$$

The fact that $\mathcal{T}^{(k)}$ is obtained from $\mathcal{T}^{(k-1)}$ by "tile composition," and $\mathcal{T}^{(k-1)}$ is obtained from $\mathcal{T}^{(k)}$ by tile subdivision is immediate from the definition of tile substitution. The formula (2.6) works for $k < 0$ in all cases. In general, we can still get the tilings $\mathcal{T}^{(k)}$ for $k > 0$ (non-uniquely), using any preimage of $\mathcal{G}^k$, which exists by surjectivity.

For $k \in \mathbb{Z}$, *prototiles of order* k are defined as $\phi^k(T) = ((\varphi^k(\mathrm{supp}(T)), i)$, where $T \in \mathcal{A}$ and i is the label of T. Tiles of order k are translates of the prototiles of order k. They are said to have "type $\mathcal{B}$" if T has type $\mathcal{B}$. The tiles of $\mathcal{T}^{(k)}$ will be called *tiles of order k obtained from $\mathcal{T}$*. The tiles of $\mathcal{T}^{(k)}$ for $k > 0$ will sometimes be referred to as "supertiles of $\mathcal{T}$".

3. Transverse Dynamics

One of the main technical ingredients in our proof will be the dynamical system $(\Omega_{\mathcal{G}}, \mathcal{G})$, restricted to a certain fractal subset defined below. Recall that $\varphi = \lambda \cdot O$, $\lambda > 1$, and O is an orthogonal matrix.

3.1. **Graph-Directed Iterated Function Systems.** Consider the directed graph $G = (V, E)$ such that the set of vertices V coincides with the alphabet $\mathcal{B}$ and the multiplicity of the set of edges from T to T' is exactly the number of occurrences of the (translate of) prototile T' in the patch $\mathcal{G}(T)$. It follows that the transpose matrix B^t is exactly the incidence matrix of the graph G. We will denote by $\mathcal{E}_{T,T'}$ the set of edges connecting a vertex T to a vertex T'. We will use the symbols $s(e), r(e)$ respectively, to denote the source and range of a directed edge.

For each vertex (prototile) $T \in V$, consider the set $S_T = \mathrm{supp}(T)$. Notice that there is a one-to-one correspondence between edges in $\mathcal{E}_{T_1,T_2}$ and the set of translates of (the order (-1) tile) $\varphi^{-1}(T_2)$ in $\varphi^{-1}(\mathcal{G}(T_1))$. *We shall fix such a correspondence.*

This can be made precise using formula (2.2) for the tile substitution: the set of edges $\mathcal{E}_{T,T'}$ corresponds to $\mathcal{D}_{T,T'}$, for $T, T' \in \mathcal{B}$. Given $e \in \mathcal{E}_{T,T'}$, we denote the corresponding vector in $\mathcal{D}_{T,T'}$ by u_e. Then the similitude

$$f_e : x \mapsto \varphi^{-1}(x + u_e) \tag{3.1}$$

maps the set $S_{T'}$ onto the translate of $\varphi^{-1}(S_{T'})$ corresponding to the edge e in the patch $\varphi^{-1}(\mathcal{G}(T))$ according to (2.3). Observe that distinct edges define different maps. Then $G = (V, E)$, $\{f_e\}_{e\in E}$, is a *graph-directed system*, and it uniquely defines a family of non-empty compact sets $\{K_T\}_{T\in\mathcal{B}}$ of $\mathbb{R}^d$ such that

$$K_T = \bigcup_{T'\in\mathcal{B}} \bigcup_{e\in\mathcal{E}_{T,T'}} f_e(K_{T'}), \tag{3.2}$$

see [MW] or [Fa, p.48]. Note that $K_T \subset S_T$ for every $T \in \mathcal{B}$. The set K_T is obtained from S_T by consecutively removing all "φ-preimages" of tiles from $\mathcal{A} \setminus \mathcal{B}$. We note that the union (3.2) need not be disjoint.

Observe that the contraction coefficient of every map f_e, $e \in E$, is exactly $1/\lambda$. Thus, to find the Hausdorff dimension of the sets $\{K_T\}_{T\in\mathcal{B}}$, one needs to consider the matrix $D^{(s)}$ with the entries

$$D^{(s)}_{T_1,T_2} = \sum_{e\in\mathcal{E}_{T_1,T_2}} \frac{1}{\lambda^s} = \frac{1}{\lambda^s} m_{T_2,T_1}.$$

It follows from [MW], see also [Fa, Corollary 3.5][1], that the Hausdorff dimension of each set K_T, $T \in \mathcal{B}$, is the unique positive number α satisfying

$$1 = \rho(D^{(\alpha)}) = \frac{1}{\lambda^\alpha}\rho(B^t) = \frac{\rho(B)}{\lambda^\alpha}.$$

Therefore, the Hausdorff dimension of every set K_T, $T \in \mathcal{B}$, is equal to

$$\alpha = \log(\rho(B))/\log(\lambda). \tag{3.3}$$

Remark 3.1. It is proved in [MW], see also [Fa, Corollary 3.5], that the α-dimensional Hausdorff measure of K_T, denoted $\mathcal{H}^\alpha(K_T)$, is positive and finite. This, together with (3.2), implies that

$$\mathcal{H}^\alpha(K_T \cap K_{T'}) = 0 \quad \text{for } T \neq T', \tag{3.4}$$

and $(\mathcal{H}^\alpha(K_T))_{T\in\mathcal{B}}$ is a left Perron-Frobenius eigenvector of B.

---

[1] This requires the *open set condition* which can be verified by setting $U_T = \mathrm{int}(S_T)$ and noting that $U_T \supset \bigcup_{T'\in\mathcal{B}} \bigcup_{e\in\mathcal{E}_{T,T'}} f_e(U_{T'})$, the union being disjoint.

We will also need the fact that $\{\mathcal{H}^\alpha|_{K_T}\}_{T\in\mathcal{B}}$ is the list of *natural self-similar graph-directed measures* on the attractors. This means that $\{\mathcal{H}^\alpha|_{K_T}\}_{T\in\mathcal{B}}$ is, up to scaling, the unique list of finite and positive Borel measures η_T on K_T, $T\in\mathcal{B}$, such that

$$\eta_T = \sum_{T'\in\mathcal{B}}\sum_{e\in\mathcal{E}_{T,T'}} \frac{1}{\rho(B)}(\eta_{T'}\circ f_e^{-1}), \tag{3.5}$$

see [E, 3.5]. Moreover, these natural measures may be obtained as projections of appropriate Markov measures on the sequence space, as we are now going to explain.

Let $T\in\mathcal{B}$. By the definition of graph-directed sets, $x\in K_T$ if and only if there is an infinite path $(e_0,e_1,\ldots)$ in the graph G such that $T=s(e_0)$ and

$$\{x\} = \bigcap_{k=0}^{\infty} K_{(e_0,\ldots,e_k)}, \tag{3.6}$$

where

$$K_{(e_0,\ldots,e_k)} = f_{e_0}\circ f_{e_1}\circ\cdots\circ f_{e_k}(K_{r(e_k)}). \tag{3.7}$$

Definition 3.2. (1) Let X_G be the two-sided edge shift space associated with the graph $G=(V,E)$, i.e.

$$X_G = \{(e_n)\in E^{\mathbb{Z}} : e_{n+1} \text{ follows } e_n \text{ in the graph } G \text{ for every } n\in\mathbb{Z}\}.$$

Formally, "e_{n+1} follows e_n in the graph G" means $r(e_{n+1})=s(e_n)$. The left shift on X_G is denoted by S.

(2) We will refer to X_G as the set of inifnite two-sided paths $(e_n)_{n\in\mathbb{Z}}$ in the graph G. We will need it in the next section; for now, let X_G^+ be the set of one-sided infinite paths $(e_n)_{n\ge 0}$ in the graph G. The *natural projection* $\pi_+ : X_G^+\to\mathbb{R}^d$ is defined by

$$\pi_+((e_n)_{n\ge0}) = \lim_{k\to\infty} f_{e_0}\circ f_{e_1}\circ\cdots\circ f_{e_k}(x_0),$$

which is independent of $x_0\in\mathbb{R}^d$. For $T\in\mathcal{B}$ let

$$X_G^+(T) = \{(e_n)_{n\ge0}\in X_G^+ :\ s(e_0)=T\} \tag{3.8}$$

be the set of infinite paths in G starting at the vertex T. Clearly, $X_G^+ = \coprod_{T\in\mathcal{B}} X_G^+(T)$ is a disjoint union. It follows from (3.6) that

$$K_T = \pi_+(X_G^+(T)).$$

Using (3.1), the natural projection π_+ can be written explicitly as follows:

$$\pi_+((e_n)_{n\ge0}) = \sum_{n=0}^{\infty}\varphi^{-n-1}u_{e_n}. \tag{3.9}$$

(3) Let $w=(w_T)_{T\in\mathcal{B}}$ be the right Perron-Frobenius eigenvector for the matrix B^t, such that $\sum_{T\in\mathcal{B}} w_T = 1$. We consider the Markov measure $\overline{\eta}$ on X_G^+, with initial probabilities (i.e. probabilities of starting at $T\in\mathcal{B}$) equal

to w_T and the probability of moving along an edge e equal to $\frac{w_{r(e)}}{\rho(B)w_{s(e)}}$. Consistency follows from the fact that w is the right eigenvector of the transition matrix for the graph, with eigenvalue $\rho(B)$. In view of Remark 3.1, we have

$$w_T = c_0^{-1}\mathcal{H}^\alpha(T) \ \text{ where } \ c_0 = \sum_{T' \in \mathcal{B}} \mathcal{H}^\alpha(T'),$$

and hence for a cylinder set $[e_0, \ldots, e_n] \subset X_G^+$ we obtain

$$\overline{\eta}([e_0, \ldots, e_n]) = \frac{w_{r(e_n)}}{\rho(B)^{n+1}} = \frac{\mathcal{H}^\alpha(r(e_n))}{c_0\rho(B)^{n+1}},. \tag{3.10}$$

The next lemma follows from the theory of graph-directed IFS (see e.g. the proof of [MW, Theorem 3]).

Lemma 3.3. *For $T \in \mathcal{B}$ consider*

$$\eta_T := \overline{\eta}|_{X_G^+(T)} \circ \pi_+^{-1},$$

that is, the natural projection of the measure $\overline{\eta}$ restricted to $X_G^+(T)$. Then $\{\eta_T\}_{T\in\mathcal{B}}$ is the list of graph-directed self-similar measures satisfying (3.5), and

$$\eta_T = c_0^{-1}\mathcal{H}^\alpha|_{K_T} \ \ \textit{for } T \in \mathcal{B},$$

where $c_0 = \sum_{T'\in\mathcal{B}} \mathcal{H}^\alpha(T')$.

3.2. "Cantorization" of tilings.

Recall that the graph-directed set K_T is defined for every prototile $T \in \mathcal{B}$. If $T' = T + x$ is a translate of a prototile $T \in \mathcal{B}$, then we write $K_{T'}$ for the set $K_T + x$.

Definition 3.4. (1) The *"cantorization"* of a tiling $\mathcal{T} \in \Omega_{\mathcal{G}}$ is the set

$$\mathcal{C}(\mathcal{T}) = \bigcup \{K_T : \ T \in \mathcal{T} \text{ and } T \text{ is type } \mathcal{B}\}.$$

(2) Denote

$$\Omega_0 = \{\mathcal{T} \in \Omega_{\mathcal{G}} : \mathbf{0} \in \mathcal{C}(\mathcal{T})\},$$

where $\mathbf{0}$ stands for the zero vector.

Next we present some equivalent conditions for the property $\mathcal{T} \in \Omega_0$, which are immediate from the definitions.

Remark 3.5. (1) We have $\mathcal{T} \in \Omega_0$ if and only if there exist $T_0 \in \mathcal{B}$ and $x \in K_{T_0}$ such that $T_0 - x \in \mathcal{T}$.

(2) We have $\mathcal{T} \in \Omega_0$ if and only if there is a nested sequence of type $\mathcal{B}$ tiles of order $-k$ obtained from $\mathcal{T}$, such that the intersection of their supports is the origin $\mathbf{0}$. More formally (compare (2.6)), we have that $\mathcal{T} \in \Omega_0$ if and only if there is a sequence of type $\mathcal{B}$ tiles $T_{-k} \in \phi^{-k}\mathcal{G}^k(\mathcal{T})$, $k \geq 1$, such that

$$\text{supp}(T_{-k-1}) \subset \text{supp}(T_{-k}), \ k \geq 0, \ \text{ and } \bigcap_{k=0}^{\infty} \text{supp}(T_{-k}) = \{\mathbf{0}\}.$$

Proposition 3.6. *(i) The set Ω_0 is compact in the tiling metric.*

(ii) The map $\mathcal{G} : \Omega_0 \to \Omega_0$ is a homeomorphism.

(iii) For every tiling $\mathcal{T} \in \Omega_0$, we have that $\mathcal{C}(\mathcal{G}^{-1}(\mathcal{T})) = \varphi^{-1}(\mathcal{C}(\mathcal{T}))$ and $\mathcal{C}(\mathcal{G}(\mathcal{T})) = \varphi(\mathcal{C}(\mathcal{T}))$.

Proof. (i) We only need to show that the set Ω_0 is closed in $\Omega_{\mathcal{G}}$. Consider a tiling $\mathcal{T} \notin \Omega_0$. Then $\mathbf{0} \notin \mathcal{C}(\mathcal{T})$. Then $\mathbf{0}$ does not belong to K_T for any tile of type $\mathcal{B}$ containing the origin, which is an open condition, since K_T is compact.

(ii) The continuity of the map $\mathcal{G} : \Omega_{\mathcal{G}} \to \Omega_{\mathcal{G}}$ is well-known and easily follows from the definition of the tiling metric. In view of Theorem 2.9, we only need to show that $\mathcal{G}(\Omega_0) = \Omega_0$ (note that tiles of type $\mathcal{B}$ belong to $\mathcal{A}_{\rm nonp}$, so $\mathcal{G}$ is one-to-one on Ω_0).

Remark 3.5(2) implies that if $\mathcal{T} \in \Omega_0$ then $\mathcal{G}(\mathcal{T}) \in \Omega_0$. Indeed, using the notation of the remark, we have $\phi T_{-k} \in \phi^{-(k-1)}\mathcal{G}^{k-1}(\mathcal{G}(\mathcal{T}))$, so $\{\phi T_{-k}\}_{k\ge 1}$ is a nested sequence of tiles of order $-(k-1)$ obtained from $\mathcal{G}(\mathcal{T})$, all of type $\mathcal{B}$, and clearly the intersection of their supports is $\{\mathbf{0}\}$.

Since the map $\mathcal{G}$ is invertible on Ω_0 by Theorem 2.9, we can find a (unique) tiling $\mathcal{T}_{-1}$ with $\mathcal{G}(\mathcal{T}_{-1}) = \mathcal{T}$. We claim that $\mathcal{T}_{-1} \in \Omega_0$. If not, then for some $k > 0$ all the tiles of order $-k$ obtained from $\mathcal{T}_{-1}$ containing the origin are of type $\mathcal{A} \setminus \mathcal{B}$. But the substitution of $\mathcal{A} \setminus \mathcal{B}$ tiles contains only $\mathcal{A} \setminus \mathcal{B}$ tiles, so we get a contradiction with the assumption that the origin lies in a $\mathcal{B}$ tile of $\mathcal{T}$. We have proved that $\mathcal{G}|_{\Omega_0}$ is a homeomorphism.

(iii) It follows from Equation (3.2) that $\phi^{-1}(\mathcal{C}(\mathcal{G}(T))) = K_T$ for every prototile $T \in \mathcal{B}$. Therefore,

$$\phi^{-1}(\mathcal{C}(\mathcal{G}(\mathcal{T}))) = \mathcal{C}(\mathcal{T}) \ \text{ for every tiling } \mathcal{T} \in \Omega_0.$$

Ths implies the last statement of the proposition. □

4. Transverse Measures

In this section we show that the dynamical system $(\Omega_0, \mathcal{G})$ is measure-theoretically isomorphic to a mixing Markov chain, namely, the edge shift on the graph G with the incidence matrix B^t, equipped with the measure of maximal entropy. The invariant measure on Ω_0 can be viewed as the conditional measure of the infinite invariant measure of the system $(\Omega_{\mathcal{G}}, \mathbb{R}^d)$.

We note that properties of $(\Omega_{\mathcal{G}}, \mathcal{G})$ as a topological dynamical system were earlier considered in [AP].

Let μ be the $\mathbb{R}^d$-ergodic measure on $\Omega_{\mathcal{G}}$ as described in Proposition 2.15. It is unique up to scaling; we will normalize it later. There exists a unique Borel σ-finite transverse measure $\mu^{\rm tr}$ on the transversal Γ such that

$$\mu(U - \Theta) = \mu^{\rm tr}(U) \cdot \mathcal{L}^d(\Theta), \tag{4.1}$$

where $\mathcal{L}^d$ is the d-dimensional Lebesgue measure and

$$U - \Theta = \{\mathcal{T} - x : \ \mathcal{T} \in U, \ x \in \Theta\},$$

for all Borel sets $U \subset \Gamma_Q$ and $\Theta \subset \mathrm{supp}(Q)$, $Q \in \mathcal{B}$, see Section 7 in [CS] for the details. (Actually, in [CS] this is only proved for U contained in a small ball centered at the origin, but the formula in stated generality follows from shift invariance of the measure μ.) This means that “locally” the measure μ behaves as a product measure.

Following [CS], we give the following definition. Recall that $\Gamma_T = \{\mathcal{T} \in \Gamma : T \in \mathcal{T}\}$.

Definition 4.1. For every $Q \in \mathcal{B}$ and $n \geq 0$, define

$$\mu^{\mathrm{tr}}_{n,Q} = \mu^{\mathrm{tr}}(\mathcal{G}^n(\Gamma_Q) - x),$$

where x is a vector such that $\mathcal{G}^n(\Gamma_Q) - x \subset \Gamma$. Since μ^{tr} is a transverse measure, the definition of $\mu^{\mathrm{tr}}_{n,Q}$ does not depend on the choice of x.

The next result follows from Lemma 5.11 in [CS] and the Perron-Frobenius theorem for primitive matrices.

Lemma 4.2. *There exists a (unique) right Perron-Frobenius eigenvector ξ for the matrix B such that*

$$\mu^{\mathrm{tr}}_{n,Q} = \frac{\xi_Q}{\rho(B)^n} \text{ for every } Q \in \mathcal{B} \text{ and } n \geq 0. \tag{4.2}$$

Let $G = (V, E)$ be the graph of the iterated function system constructed in Section 3.1.

Definition 4.3. (1) The *itinerary* of $\mathcal{T} \in \Omega_0$ for the $\mathcal{G}$-dynamics is a two-sided infinite path $\beta(\mathcal{T}) = (e_n)_{n\in\mathbb{Z}} \in X_G$ (see Definition 3.2), defined as follows: $\beta(\mathcal{T}) = (e_n)_{n\in\mathbb{Z}}$ if for all $n \in \mathbb{Z}$ the tiling $\mathcal{G}^n(\mathcal{T})$ has a tile T_n of type $s(e_n) = r(e_{n-1}) \in \mathcal{B}$ containing the origin and T_{n+1} occurs in $\mathcal{G}(T_n)$ in the position corresponding to e_n.

(2) Observe that $\phi^n T_{-n}$ for $n > 0$ forms an increasing sequence of supertiles of the tiling $\mathcal{T}$. We will call it the *compatible sequence of supertiles of $\mathcal{T}$ containing the origin.*

(3) Note that the itinerary need not be unique. Denote by Ω_0^* the set of all tilings $\mathcal{T} \in \Omega_0$ for which the itinerary is unique. The itinerary is non-unique if and only if for some $n \in \mathbb{Z}$, the origin $\mathbf{0}$ lies on the common boundary of two tiles of $\mathcal{B}$ type $T_n, T_n' \in \mathcal{G}^n(\mathcal{T})$ and, moreover, $\mathbf{0} \in K_{T_n} \cap K_{T_n'}$. Note that just being on the boundary of a tile may not lead to non-uniqueness.

Thus $\beta : \Omega_0^* \to X_G$ is a well-defined function, whereas β may be considered as a multi-valued function on all of Ω_0. Observe that Ω_0^* is a $\mathcal{G}$-invariant Borel subset of Ω_0.

(4) By definition, $\beta \circ \mathcal{G} = S \circ \beta$, where S is the left shift on X_G. This holds, in an appropriate sense, even when the itinerary is non-unique.

Remark 4.4. (1) Many properties of the tiling dynamical system can be expressed using the symbolic dynamics provided by the itinerary β. In particular, if we fix the left one-sided sequence $(e_n)_{n<0}$, this corresponds to the set of translates of $\mathcal{T} \in \Omega_0^*$, such that the origin stays in the interior of its original tile of type $r(e_{-1})$. This is a "piece" of the translation orbit of $\mathcal{T}$. On the other hand, fixing the right half of the symbolic orbit $(e_n)_{n\ge 0}$ corresponds to the transversal; more precisely, for all $\mathcal{T} \in (\Gamma_T + x) \cap \Omega_0^*$ for a fixed vector x, the sequences $\beta(\mathcal{T})$ agree in $n \ge 0$.

(2) There are, however, some complications. First, β is not well-defined on $\Omega_0 \setminus \Omega_0^*$. Second, β need not be one-to-one and need not be onto (even if extended to Ω_0 as a multi-valued function). The reason is that the sequence $(e_n)_{n<0}$ determines a sequence of compatible supertiles *whose union need not be the entire space* $\mathbb{R}^d$. We will deal with such sequences by showing that they have zero measure of maximal entropy for S.

Definition 4.5. (1) Define X_G^* as the set of $(e_n) \in X_G$ such that the compatible increasing sequence of supertiles, corresponding to $(e_n)_{n<0}$, has all of $\mathbb{R}^d$ as the limit (i.e. the union) of the supports. It is clear that X_G^* is S-invariant.

(2) We define the *natural projection map* $\pi : X_G^* \to \Omega_0$ so that (e_n) is an itinerary of $\mathcal{T} := \pi(\overline{e})$. It is possible to describe $\mathcal{T}$ explicitly, as a limit of an increasing compatible sequence of patches (whose supports are the supports of supertiles of $\mathcal{T}$). The condition $\overline{e} = (e_n)_{n\in\mathbb{Z}} \in X_G$ means, by definition, that

$$r(e_n) + u_{e_n} \in \mathcal{G}(s(e_n)) \quad \text{for all } n \in \mathbb{Z} \tag{4.3}$$

(recall that the vertices of G are identified with the prototiles in $\mathcal{B}$). A tile of $\mathcal{T} = \pi(\overline{e})$ containing the origin (possibly non-unique) must be

$$T_0 - \sum_{n=0}^{\infty} \varphi^{-n-1} u_{e_n} = T_0 - \pi_+(\overline{e}_+), \tag{4.4}$$

where $T_0 = s(e_0)$ and $\overline{e}_+ = (e_n)_{n\ge 0}$ (recall that π_+ was defined in Definition 3.2(2)). Note that this already guarantees $\mathcal{T} \in \Omega_0$, in view of Remark 3.5(1) and (3.9). Now we let

$$\pi(\overline{e}) = \lim_{k\to\infty} \left[\mathcal{G}^k(s(e_{-k})) - \sum_{n=-k}^{\infty} \phi^{-n-1} u_{e_n} \right]. \tag{4.5}$$

We claim that these patches are increasing and compatible. Indeed,

$$\mathcal{G}^k(s(e_{-k})) - \sum_{n=-k}^{\infty} \phi^{-n-1} u_{e_n} \subset \mathcal{G}^{k+1}(s(e_{-k-1})) - \sum_{n=-k-1}^{\infty} \phi^{-n-1} u_{e_n}$$

reduces to

$$u_{e_{-k-1}} + s(e_{-k}) \in \mathcal{G}(s(e_{-k-1})),$$

which follows from (4.3), keeping in mind that $s(e_{-k}) = r(e_{-k-1})$. Thus, the right-hand side of (4.5) is well-defined, and it is a tiling of the entire $\mathbb{R}^d$ if $\overline{e} \in X_G^*$.

Lemma 4.6. *We have $\pi \circ S = \mathcal{G} \circ \pi$ on X_G^* and $\pi \circ \beta = Id$ on $\beta(\Omega_0^*)$.*

Proof. This is an immediate consequence of the definitions. □

Next we consider the measure of maximal entropy (the Parry measure) $\overline{\nu}$ for the shift S on X_G. Recall that the incidence matrix for the graph G is B^t. The Parry measure (of the edge shift) is a Markov measure, given by

$$\overline{\nu}([e_k,\dots,e_{n+k}]) = u_{s(e_k)}v_{s(e_k)}\prod_{k=0}^{n}\frac{v_{r(e_{n+k})}}{\rho(B)v_{s(e_{n+k})}} = \frac{u_{s(e_k)}v_{r(e_{n+k})}}{\rho(B)^{n+1}}, \tag{4.6}$$

where $[e_k,\dots,e_{n+k}]$ is a cylinder set in X_G, for any $k \in \mathbb{Z}$ and $n \geq 0$, $u = (u_Q)_{Q\in\mathcal{B}}$ is the left Perron-Frobenius eigenvector of B^t, and $v = (v_Q)_{Q\in\mathcal{B}}$ is the right Perron-Frobenius eigenvector of B^t, normalized so that $\sum_{Q\in\mathcal{B}} u_Q v_Q = 1$. The measure is clearly shift-invariant. We have

$$\nu(\{\overline{e}\in X_G:\ s(e_0)=Q\}) = \sum_{Q'\in\mathcal{B}}\sum_{e\in\mathcal{E}_{Q,Q'}} u_Q v_{Q'}\cdot\rho(B)^{-1} = u_Q v_Q,$$

which implies that $\overline{\nu}$ is a probability measure. For the vector u we can take the vector ξ from Lemma 4.2, which is a right Perron-Frobenius eigenvector for B, and for the vector v we can take $(\mathcal{H}^\alpha(K_Q))_{Q\in\mathcal{B}}$, which is a left Perron-Frobenius eigenvector for B, see Remark 3.1. Since the measure μ was defined up to scaling, we can normalize it (this will also affect the transverse measure) so that

$$\sum_{Q\in\mathcal{B}}\xi_Q\mathcal{H}^\alpha(K_Q) = \sum_{Q\in\mathcal{B}}\mu^{\mathrm{tr}}(\Gamma_Q)\mathcal{H}^\alpha(K_Q) = 1. \tag{4.7}$$

Then we have

$$\overline{\nu}([e_k,\dots,e_{k+n}]) = \frac{\xi_{s(e_k)}}{\rho(B)^{n+1}}\,\mathcal{H}^\alpha(K_{r(e_{k+n})})\ \text{ for } k\in\mathbb{Z},\ n\geq 0. \tag{4.8}$$

It is well-known that the measure-preserving transformation $(X_G, S, \overline{\nu})$ is ergodic, where S is the left shift.

Lemma 4.7. *We have* $\overline{\nu}(X_G\setminus X_G^*) = 0$.

Proof. Recall that there is an integer $k > 0$ such that for each prototile $T\in\mathcal{B}$ the interior of the patch $\mathcal{G}^k(T)$ contains a translate of T. We can assume without loss of generality, passing from $\mathcal{G}$ to $\mathcal{G}^k$, that $k = 1$. Then for any vertex of the graph G (i.e. $Q\in\mathcal{B}$) there is an edge e, with $s(e) = Q$, which corresponds to the choice of an interior tile in the patch $\mathcal{G}(Q)$. A one-sided path $(e_n)_{n<0}$, which includes infinitely many of these "interior" edges, will necessarily belong to X_G^*. Indeed, choosing an interior supertile of order n inside the supertile of order $n+1$, for $n > 0$, implies that the union of the compatible sequence of supertiles contains the ball of raduis $\delta\lambda^n$ centered at the origin, for some $\delta > 0$. A standard argument shows that this is a full measure set. To verify this, note that the set of paths, which avoid the selected edges, has a growth rate equal to the spectral radius of a matrix B' having at least one entry in each row smaller than that of B, whence $\rho(B') < \rho(B)$. □

Definition 4.8. Define the measure ν on Ω_0 as the "push-forward" of $\overline{\nu}$ on X_G^* via the map π, that is,

$$\nu(U) = \overline{\nu}(\pi^{-1}(U)) \quad \text{for Borel } U \subset \Omega_0.$$

Since $\overline{\nu}$ is S-invariant, we obtain from Lemma 4.6 that the measure ν is $\mathcal{G}$-invariant on Ω_0.

Lemma 4.9. *We have* $\nu(\Omega_0 \setminus \Omega_0^*) = 0$.

Proof. The argument is almost the same as in the proof of Lemma 4.7. It is enough to prove that the set of tilings $\mathcal{T} \in \Omega_0$, for which there exists $n \in \mathbb{Z}$ such that $\mathbf{0}$ is on the boundary of a tile of type $\mathcal{B}$ in $\mathcal{G}^n(\mathcal{T})$, has ν measure zero. Considering the itineraries of such tilings, we see that they must contain only finitely many edges e_i corresponding to the tile of type $r(e_i)$ in the interior of $\mathcal{G}(s(e_i))$, for $i \geq 0$. But the growh rate of such sequences is strictly less than $\rho(B)$, hence their $\overline{\nu}$ measure equals zero, as desired. $\square$

Theorem 4.10. *Suppose that the Markov measure* $\overline{\nu}$ *is defined by (4.8), using the normalization (4.7), and* $\nu = \overline{\nu} \circ \pi^{-1}$. *Then the following hold:*

(i) The probability-preserving system $(\Omega_0, \mathcal{G}, \nu)$ *is measure-theoretically isomorphic to* $(X_G, S, \overline{\nu})$, *hence ergodic.*

(ii) For any $Q \in \mathcal{B}$ *and all Borel sets* $\Theta \subset \Gamma_Q$, $W \subset K_Q$ *we have*

$$\nu(\Theta - W) = \mu^{\mathrm{tr}}(\Theta) \cdot \mathcal{H}^{\alpha}(W). \tag{4.9}$$

Proof. (i) This follows from Lemmas 4.6, 4.7, and 4.9.

(ii) The left-hand side of (4.9) is well-defined, since $\Gamma_Q - K_Q \subset \Omega_0$ by Remark 3.5(1). First let us prove the equality for $\Theta = \Gamma_Q$. Recall that $X_G^+(Q)$ denotes the set of one-sided paths in G starting at Q. We have

$$\nu(\Gamma_Q - W) = \overline{\nu}\left(\{\overline{e} \in X_G : \ \overline{e}_+ \in X_G^+(Q) \text{ and } \pi_+(\overline{e}_+) \in W\}\right),$$

using the fact that $\mathcal{T} \in \Gamma_Q - W$, with $W \subset K_Q$, has an itinerary with $s(e_0) = Q$, and ν almost every tiling has a unique itinerary by Lemma 4.9. The measure $\overline{\nu}$ on X_G induces a measure $\overline{\nu}_+$ on X_G^+ via the projection $\overline{e} \mapsto \overline{e}_+$. Comparing (4.6) with (3.10) we see that

$$\overline{\nu}_+|_{X_G^+(Q)} = c_0^{-1}\xi_Q \cdot \overline{\eta}|_{X_G^+(Q)}.$$

Thus,

$$\begin{aligned} \nu(\Gamma_Q - W) &= \overline{\nu}_+|_{X_G^+(Q)}(\pi_+^{-1}W) \\ &= c_0^{-1}\xi_Q \cdot \overline{\eta}|_{X_G^+(Q)}(\pi_+^{-1}W) \\ &= c_0^{-1}\xi_Q \cdot \eta_Q(W) \\ &= \xi_Q \mathcal{H}^{\alpha}(W), \end{aligned}$$

where we used Lemma 3.3 in the last step.

Now let us verify (4.9) for an arbitrary Borel $\Theta \subset \Gamma_Q$. The transversal Γ_Q is topologically a Cantor set, in which the Borel σ-algebra is generated by the sets of the form $\mathcal{G}^n(\Gamma_{Q'}-x)$, $Q' \in \mathcal{A}$, where x is such that $Q+x \in \mathcal{G}^n(Q')$. We have

$$\nu(\mathcal{G}^n(\Gamma_{Q'} - x - W) = \nu(\mathcal{G}^n(\Gamma_{Q'} - \phi^{-n}(x+W))) = \nu(\Gamma_{Q'} - \phi^{-n}(x+W)),$$

using the fact that ν is $\mathcal{G}$-invariant. Note that $W \subset K_Q$ and $Q+x \in \mathcal{G}^n(Q')$ imply $W + x \subset \mathcal{C}(\mathcal{G}^n(Q')) = \phi^n K_{Q'}$, hence $\phi^{-n}(x+W) \subset K_{Q'}$, and by the case of (4.9) already proved,

$$\begin{aligned} \nu(\Gamma_{Q'} - \phi^{-n}(x+W)) &= \xi_{Q'} \cdot \mathcal{H}^\alpha(\phi^{-n}W) \\ &= \frac{\xi_{Q'} \cdot \mathcal{H}^\alpha(W)}{\lambda^{nd}} \\ &= \frac{\xi_{Q'} \cdot \mathcal{H}^\alpha(W)}{\rho(B)^{n+1}} \\ &= \mu^{\mathrm{tr}}(\Gamma_{Q'} - x) \cdot \mathcal{H}^\alpha(W), \end{aligned}$$

by Lemma 4.2 and Definition 4.1. The proof is complete.

□

5. Second Order Ergodic Theorem

In this section we establish the second order theorem for tiling substitution systems. We begin with Lemma 5.3 saying that the second order ergodic theorem can be established by checking the convergence of second order averages for one (any) function only. This lemma was originally proved in [Fi1, Theorem 4] for the discrete case. We include the proof for the reader's convenience. The proof is based on the following generalization of Hopf's ratio ergodic theorem. Recall that a group action is *free* if identity is the only group element for which there exists a fixed point. Our tiling translation action is free in the measure-theoretic sense, since tilings containing at least one tile of type $\mathcal{B}$ are non-periodic, and these tilings form an invariant set of full μ measure. Recall that B_R denotes the closed Euclidean ball.

Theorem 5.1 (M. Hochman [H]). *Let $\{T^u\}_{u\in\mathbb{R}^d}$ be a free ergodic measure preserving action on a standard σ-finite measure space X. Then for μ-a.e. $x \in X$ and every $f, g \in L^1(X,\mu)$ with $\int_X g d\mu \neq 0$, we have*

$$\frac{\int_{B_R} f(T^u(x))du}{\int_{B_R} g(T^u(x))du} \to \frac{\int_X f d\mu}{\int_X g d\mu} \quad \textit{as } R \to \infty.$$

Remark 5.2. In fact, [H] considers non-singular free ergodic actions of $\mathbb{Z}^d$ or $\mathbb{R}^d$, which includes measure-preserving actions, and averaging is over balls in any norm. We note that for our purposes it would suffice to use an older ratio ergodic theorem of M. Becker [Beck], but it would require a little additional argument, so we chose to quote the recent more general result of M. Hochman.

Lemma 5.3. *Let $\{T^u\}_{u\in\mathbb{R}^d}$ be a free ergodic measure preserving action on a standard σ-finite measure space (X,μ). Assume that there exists $\alpha > d-1$ such that for some function $g \in L^\infty(X,\mu)$ with $\int_X g\,d\mu \neq 0$, the limit*

$$\lim_{t\to\infty} \frac{1}{\log(t)} \int_0^t \frac{\int_{B_R} g(T^u(x))\,du}{R^{\alpha+1}}\,dR$$

exists and is finite for μ a.e. $x \in X$. Then

(i) this limit is constant almost everywhere;

(ii) writing this limit as $c \cdot \int_X g d\mu$ for some constant c, we get that for every function $f \in L^1(X,\mu)$,

$$\lim_{t\to\infty} \frac{1}{\log(t)} \int_0^t \frac{\int_{B_R} f(T^u(x))\,du}{cR^{\alpha+1}}\,dR = \int_X f\,d\mu$$

for μ-a.e. $x \in X$.

Proof. (i) Set

$$S_R^g(x) = \int_{B_R} g(T^u(x))\,du \quad \text{and} \quad \overline{g}(x) = \lim_{t\to\infty} \frac{1}{\log(t)} \int_0^t \frac{S_R^g(x)}{R^{\alpha+1}}\,dR.$$

By assumption, the function $\overline{g}(x)$ is finite for μ-a.e. $x \in X$, and it is straightforward to check that $\overline{g}$ is measurable. We claim that $\overline{g}(T^v(x)) = \overline{g}(x)$ for every $v \in \mathbb{R}^d$ and μ-a.e. $x \in X$. Indeed,

$$\int_{B_R} g(T^u(x))\,du - \int_{B_R} g(T^{u+v}(x))\,du = O(R^{d-1}\|g\|_\infty) \quad \text{as } R \to \infty,$$

and the assumption $\alpha > d-1$ yields the claim. Since the measure μ is ergodic, we get that the function $\overline{g}$ is constant almost everywhere.

(ii) Applying Theorem 5.1, we obtain that

$$\frac{S_R^f(x)}{S_R^g(x)} \to \frac{\int f\,d\mu}{\int g\,d\mu} \quad \text{as } R \to \infty.$$

Assume for definiteness that $\int g\,d\mu > 0$ and $\int f\,d\mu \geq 0$. It follows that for any $\varepsilon > 0$ there is $R_0 > 0$ such that for all $R > R_0$,

$$S_R^g(x)\frac{\int f\,d\mu}{\int g\,d\mu}(1-\varepsilon) \leq S_R^f(x) \leq S_R^g(x)\frac{\int f d\mu}{\int g\,d\mu}(1+\varepsilon).$$

Dividing the inequalities by $R^{\alpha+1}$ and integrating with respect to R yields

$$(1-\varepsilon)\int g\,d\mu\,\frac{\int f\,d\mu}{\int g\,d\mu} \leq \lim_{t\to\infty} \frac{1}{\log(t)} \int_0^t \frac{S_R^f(x)}{R^{d+1}}\,dR \leq (1+\varepsilon)\int g\,d\mu\,\frac{\int f\,d\mu}{\int g\,d\mu}.$$

Taking the limit as $\varepsilon \to 0$, we obtain the result. □

Now we are ready to prove the main result of the paper.

Theorem 5.4. *Let $\mathbb{X} = (\Omega_{\mathcal{G}}, \mu, \mathbb{R}^d)$ be the tiling dynamical system corresponding to a tile substitution $\mathcal{G}$. Suppose that the tiling system satisfies the assumptions of Section 2.3. Assume that μ is an infinite (σ-finite) invariant measure, positive and finite on $\Omega_{\mathcal{B}}$, where $\Omega_{\mathcal{B}}$ is the set of tilings which has a type $\mathcal{B}$ tile covering the origin.*

Then there exist positive parameters α and c such that for μ-almost every tiling $\mathcal{T} \in \Omega$ and for every function $f \in L^1(\Omega, \mu)$, we have

$$\lim_{t\to\infty} \frac{1}{\log(t)} \int_0^t \frac{\int_{B_R} f(\mathcal{T} - u)\,du}{cR^\alpha} \frac{dR}{R} = \int_{\Omega_{\mathcal{G}}} f d\mu. \tag{5.1}$$

Here $\alpha = \log(\rho(B))/\log(\lambda)$ is the Hausdorff dimension of the graph-directed self-similar sets from Section 3.1 and

$$c = \gamma \cdot \lim_{k\to\infty} \frac{1}{k} \int_0^k \frac{\mathcal{H}^\alpha(B_{\lambda^{-t}}(u) \cap K_Q)}{\lambda^{-t\alpha}} dt$$

for $\mathcal{H}^\alpha$-a.e. $u \in K_Q$ and for every $Q \in \mathcal{B}$, where $\mathcal{H}^\alpha$ is the α-dimensional Hausdorff measure on K_Q . The parameter c is the average density of K_Q, and γ is a normalizing constant:

$$\gamma^{-1} = \sum_{Q\in\mathcal{B}} \xi_Q \mathcal{H}^\alpha(Q), \quad \textit{where} \quad \sum_{Q\in\mathcal{B}} \xi_Q \mathcal{L}^d(Q) = \mu(\Omega_{\mathcal{B}}),$$

and $(\xi_Q)_{Q\in\mathcal{B}}$ is a right Perron-Frobenius eigenvector of the matrix B.

Proof. (1) First note that, without loss of generality, we can normalize μ in such a way that (4.7) holds, so that $\gamma = 1$. We then define ν as in Theorem 4.10 and consider the ergodic probability-preserving transformation $(\Omega_0, \nu, \mathcal{G}^{-1})$. We follow, in part, the argument of [Fa, Theorem 6.6]. Recall that B_R denotes the closed ball of radius R centered at the origin. Define a function $\psi : \Omega_0 \to \mathbb{R}$ by

$$\psi(\mathcal{T}) = \int_0^1 \frac{\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T}) \cap B_{\lambda^t})}{\lambda^{t\alpha}} dt.$$

Since the Hausdorff measure $\mathcal{H}^\alpha$ is finite on sets $\mathcal{K}_T$, $T \in \mathcal{B}$, and the ball B_λ contains only a finite number of tiles, the function ψ is bounded. It is straightforward to check that the function ψ is measurable.

(2) Recall that $\varphi = \lambda \cdot O$, where O is an orthogonal matrix, hence $\mathcal{H}^\alpha(\phi^{-1}E) = \lambda^{-d}\mathcal{H}^\alpha(E)$ for any Borel set E. Note also that $\varphi(B_{\lambda^t}) = B_{\lambda^{t+1}}$.

Applying Proposition 3.6, we obtain that

$$\begin{aligned}\psi(\mathcal{G}^{-1}(\mathcal{T})) &= \int_0^1 \frac{\mathcal{H}^\alpha(\varphi^{-1}(C(\mathcal{T}))\cap B_{\lambda^t})}{\lambda^{t\alpha}}\,dt\\ &= \int_0^1 \frac{\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T})\cap B_{\lambda^{t+1}})}{\lambda^{(t+1)\alpha}}\,dt\\ &= \int_1^2 \frac{\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T})\cap B_{\lambda^t})}{\lambda^{t\alpha}}\,dt.\end{aligned}$$

It follows that

$$\sum_{i=0}^{k-1}\psi(\mathcal{G}^{-i}\mathcal{T}) = \int_0^k \frac{\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T})\cap B_{\lambda^t})}{\lambda^{t\alpha}}\,dt.$$

Thus, applying the Birkhoff Ergodic Theorem to the system $(\Omega_0,\nu,\mathcal{G}^{-1})$ and the function ψ, we get that

$$\lim_{k\to\infty}\frac{1}{k}\int_0^k \frac{\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T})\cap B_{\lambda^t})}{\lambda^{t\alpha}}\,dt = \int_{\Omega_0}\psi(\mathcal{S})\,d\nu(\mathcal{S}) \tag{5.2}$$

for ν-a.e. tiling $\mathcal{T}\in\Omega_0$. Substituting $\lambda^t = R$ into (5.2), we obtain that

$$\int_{\Omega_0}\psi(\mathcal{S})\,d\nu(\mathcal{S}) = \lim_{z\to\infty}\frac{1}{\log(z)}\int_1^z \frac{\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T})\cap B_R)}{R^\alpha}\,\frac{dR}{R} \tag{5.3}$$

for ν-a.e. $\mathcal{T}\in\Omega_0$. (Passing from $z=\lambda^k$ for $k\in\mathbb{N}$ to an arbitrary $z>0$, $z\to\infty$, in the limit above is justified, since ψ is a bounded function.)

(3) We have

$$\Omega_{\mathcal{G}} = \bigcup_{Q\in\mathcal{A}}(\Gamma_Q - \mathrm{supp}(Q)).$$

The sets in the right-hand side have intersections of zero μ measure in view of (4.1), since $\mathcal{L}^d(\partial(\mathrm{supp}(Q)))=0$. Consider the function

$$g := \sum_{Q\in\mathcal{B}}\frac{\mathcal{H}^\alpha(Q)}{\mathcal{L}^d(Q)}\cdot\chi_{\Gamma_Q-\mathrm{supp}(Q)} \in L^\infty(\Omega_{\mathcal{G}},\mu).$$

That is, $g(\mathcal{T})$ is nonzero if and only if the origin lies in a $\mathcal{T}$-tile $Q-x$ of type $\mathcal{B}$, and then the value of the function is $\frac{\mathcal{H}^\alpha(Q)}{\mathcal{L}^d(Q)}$ (this is well-defined on a set of full μ measure). Then (4.1) implies

$$\int_{\Omega_{\mathcal{G}}} g(\mathcal{T})\,d\mu(\mathcal{T}) = \sum_{Q\in\mathcal{B}}\mu^{\mathrm{tr}}(\Gamma_Q)\mathcal{H}^\alpha(Q) = 1.$$

In view of Lemma 5.3, it suffices to establish the second order ergodic theorem just for the function g.

Given a tiling $\mathcal{T} \in \Omega_{\mathcal{G}}$, denote

$$V_R(\mathcal{T}) = \int_{B_R} g(\mathcal{T} - u)\, du.$$

Observe that for every tile $T = Q - x \in \mathcal{T}$, with $Q \in \mathcal{B}$, such that $\mathrm{supp}(T) \subset B_R$, integrating $g(\mathcal{T} - u)$ over $\mathrm{supp}(T)$ contributes $\mathcal{H}^\alpha(K_T)$ to $V_R(\mathcal{T})$. Exactly the same contribution from T comes to $\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T}) \cap B_R)$. Therefore, the difference between $\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T}) \cap B_R)$ and $V_R(\mathcal{T})$ is bounded (in modulus) by the sum of $\mathcal{H}^\alpha(K_T)$ over those $T \in \mathcal{T}$ of type $\mathcal{B}$ whose supports intersect ∂B_R. Thus, denoting by d_M the maximal diameter of a prototile, we obtain

$$|\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T}) \cap B_R) - V_R(\mathcal{T})| \le \mathcal{L}^d(B_R \setminus B_{R-d_M}) \cdot \max_{Q \in \mathcal{B}} \frac{\mathcal{H}^\alpha(Q)}{\mathcal{L}^d(Q)} = O(R^{d-1}),$$

with the implied constant in $O(\cdot)$ depending only on the tiling $\mathcal{T}$.

Since $\alpha > d - 1$ (one of our standing assumptions), we obtain that

$$\lim_{z \to \infty} \frac{1}{\log(z)} \int\limits_1^z \frac{|\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T}) \cap B_R) - V_R(\mathcal{T})|}{R^{\alpha+1}}\, dR = 0.$$

Using Equation (5.3), we conclude that

$$\lim_{z \to \infty} \frac{1}{\log(z)} \int\limits_1^z \frac{\int_{B_R} g(\mathcal{T} - u) du}{R^{\alpha+1}}\, dR = \int_{\Omega_0} \psi(\mathcal{S})\, d\nu(\mathcal{S}) \tag{5.4}$$

for ν-a.e. tiling $\mathcal{T} \in \Omega_0$. Denote by Y the set of all tilings in $\Omega_{\mathcal{G}}$ for which Equation (5.4) holds. Observe that if $\mathcal{T} \in Y$, then $\mathcal{T} - v \in Y$ for any $v \in \mathbb{R}^d$ (here we use $\alpha > d - 1$ again), i.e. Y is translation-invariant. Translation invariance of Y implies that $Y \cap (\Gamma_Q - K_Q) = (Y \cap \Gamma_Q) - K_Q$ for each prototile $Q \in \mathcal{B}$. Since $\nu(Y \cap \Omega_0) > 0$, there is a prototile $Q \in \mathcal{B}$ with $\nu(Y \cap (\Gamma_Q - K_Q)) > 0$. Then Theorem 4.10(ii) implies that

$$0 < \nu(Y \cap (\Gamma_Q - K_Q)) = \nu((Y \cap \Gamma_Q) - K_Q) = \mu^{\mathrm{tr}}(Y \cap \Gamma_Q) \cdot \mathcal{H}^\alpha(K_Q),$$

hence $\mu^{\mathrm{tr}}(Y \cap \Gamma_Q) > 0$. Again using translation-invariance of Y and (4.1) we obtain

$$\mu(Y) \ge \mu((Y \cap \Gamma_Q) - \mathrm{supp}(Q)) = \mu^{\mathrm{tr}}(Y \cap \Gamma_Q) \cdot \mathcal{L}^d(\mathrm{supp}(Q)) > 0,$$

and ergodicity of the tiling dynamical system $\Omega_{\mathcal{G}}, \mu, \mathbb{R}^d)$ implies that (5.4) holds for μ-a.e. $\mathcal{T} \in \Omega_{\mathcal{G}}$. Setting

$$c = \int_{\Omega_0} \psi(\mathcal{S})\, d\nu(\mathcal{S}),$$

we get the result.

(4) It remains to show that the parameter c can be interpreted as the average density of the graph-directed set. Using the same arguments as in (2) above, we obtain that

$$\begin{aligned}\psi(\mathcal{G}^k(\mathcal{T})) &= \int_0^1 \frac{\mathcal{H}^\alpha(\varphi^k(\mathcal{C}(\mathcal{T}))\cap B_{\lambda^t})}{\lambda^{t\alpha}}\,dt \\ &= \int_0^1 \frac{\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T})\cap B_{\lambda^{t-k}})}{\lambda^{(t-k)\alpha}}\,dt \\ &= \int_{-k}^{-k+1} \frac{\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T})\cap B_{\lambda^t})}{\lambda^{t\alpha}}\,dt.\end{aligned}$$

Applying the Birkhoff Ergodic Theorem to the system $(\Omega_0,\nu,\mathcal{G})$ and the function ψ, we see that for ν-a.e. $\mathcal{T}\in\Omega_0$,
(5.5)

$$\lim_{k\to\infty}\frac{1}{k}\int_{-k}^{0}\frac{\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T})\cap B_{\lambda^t})}{\lambda^{t\alpha}}\,dt = \lim_{k\to\infty}\frac{1}{k}\int_{0}^{k}\frac{\mathcal{H}^\alpha(\mathcal{C}(\mathcal{T})\cap B_{\lambda^{-t}})}{\lambda^{-t\alpha}}\,dt = c.$$

We have

$$\Omega_0 = \bigcup_{Q\in\mathcal{B}}(\Gamma_Q - K_Q),$$

and the sets in the right-hand side have intersections of zero ν measure, in view of Theorem 4.10(ii) and (3.4). Then the set of tilings $\mathcal{T}\in\Omega_0$, such that (5.5) holds and $\mathcal{T}$ belongs to $\Gamma_Q - \mathcal{K}_Q$ for a unique $Q\in\mathcal{B}$, has full ν measure. Denote this set of tilings by Z. Observe that the behavior of the limit in (5.5) depends only on the small neighborhood of the origin (because the limit doesn't change if we replace $\int_0^k$ by $\int_i^k$ for any fixed $i\in\mathbb{N}$), hence $\mathcal{T}-u\in Z$ for $T\in\Gamma_Q$ and $u\in K_Q$ implies $\mathcal{T}'-u\in Z$ for any $\mathcal{T}'\in\Gamma_Q$. Thus, for every $Q\in\mathcal{B}$,

$$Z\cap(\Gamma_Q - K_Q) = \Gamma_Q - K_Q'$$

for some $K_Q'\subset K_Q$. We have

$$\begin{aligned}\mu^{\rm tr}(\Gamma_Q)\cdot\mathcal{H}^\alpha(K_Q) &= \nu(\Gamma_Q - K_Q) \\ &= \nu(Z\cap(\Gamma_Q - K_Q)) \\ &= \nu(\Gamma_Q - K_Q') \\ &= \mu^{\rm tr}(\Gamma_Q)\cdot\mathcal{H}^\alpha(K_Q').\end{aligned}$$

It follows that $\mathcal{H}^\alpha$-a.e. $u\in K_Q$ is such that $\mathcal{T}-u\in Z$ for $T\in\Gamma_Q$, which means, rewriting (5.5), that

$$c = \lim_{k\to\infty}\frac{1}{k}\int_0^k\frac{\mathcal{H}^\alpha(K_Q\cap B_{\lambda^{-t}}(u))}{\lambda^{-t\alpha}}\,dt,$$

as desired. The proof is complete. □

Remark 5.5. 1. Since there exists $f \in L^1(\Omega_{\mathcal{G}}, \mu)$ such that $0 < \int_{\Omega_{\mathcal{G}}} f dm < \infty$, it is immediate that the paramaters α and c in Theorem 5.4 are invariants of measure-theoretic isomorphism.

2. We considered averaging over Euclidean balls in Theorem 5.4. This was needed in the equality $\phi(B_R) = B_{\lambda R}$. If we restrict ourselves to the case when ϕ is a pure dilation, i.e. $\phi(x) = \lambda x$ for $\lambda > 1$, then we can use averaging over balls in any norm.

6. Substitution Dynamical Systems

In this section, we derive the second order ergodic theorem for a class of one-dimensional substitution systems. We begin with a brief review of the background. This will be reminiscent of our discussion of the structure of tiling substitutions, however, there are certain fundamental differences. One of the principal differences between symbolic substitution systems and their tiling counterparts is that symbolic substitutions may have finite ergodic invariant measures supported off the minimal components.

Now $\mathcal{A}$ is a finite alphabet (usually $\mathcal{A} = \{1, \dots, N\}$) and $\mathcal{A}^+$ is the set of all finite non-empty words over $\mathcal{A}$. A map $\sigma : \mathcal{A} \to \mathcal{A}^+$ is called a *substitution*; it is extended to $\mathcal{A}^+$ by concatenation. Given two words $v, w \in \mathcal{A}^+$, we will write $v \prec w$ if v is a subword of w. Denote by $L(\sigma)$ the set of all words $w \in \mathcal{A}^+$ such that $w \prec \sigma^n(a)$ for some $a \in \mathcal{A}$ and $n \geq 1$. The family $L(\sigma)$ is called the *language* of the substitution. For a word w, its length will be denoted by $|w|$.

Definition 6.1. The *substitution dynamical system* determined by a substitution σ is a pair (X_σ, S), where

$$X_\sigma = \{x \in \mathcal{A}^{\mathbb{Z}} : \ x[-n, n] \in L(\sigma) \text{ for all } n \geq 1\}$$

and $S : \mathcal{A}^{\mathbb{Z}} \to \mathcal{A}^{\mathbb{Z}}$ is the left shift. The set X_σ is S-invariant and closed in $\mathcal{A}^{\mathbb{Z}}$ with respect to the product topology.

Given $a, b \in \mathcal{A}$, denote by $m_{a,b}$ the number of occurrences of a in the word $\sigma(b)$. The matrix $M(\sigma) = (m_{a,b})_{a,b \in \mathcal{A}}$ is the *matrix of the substitution σ*. Reordering the letters in the alphabet $\mathcal{A}$, the matrix $M(\sigma)$ can be transformed to have an upper block-triangular form:

$$M_\sigma = \begin{pmatrix} F_1 & 0 & \cdots & 0 & X_{1,s+1} & \cdots & X_{1,m} \\ 0 & F_2 & \cdots & 0 & X_{2,s+1} & \cdots & X_{2,m} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & \cdots & F_s & X_{s,s+1} & \cdots & X_{s,m} \\ 0 & 0 & \cdots & 0 & F_{s+1} & \cdots & X_{s+1,m} \\ \vdots & \vdots & \cdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & 0 & \cdots & F_m \end{pmatrix} \tag{6.1}$$

where diagonal matrices F_i are irreducible, and for any fixed $j = s+1, ..., m$, at least one of the matrices $X_{k,j}$ is non-zero. Replacing σ by σ^k if needed, we can ensure that all matrices F_i are zero or primitive.

Denote by $\mathcal{A}_i$ the set of letters corresponding to the matrix F_i, $i = 1, \ldots, m$. For $i = 1, \ldots, m$, denote by Y_i the set of all $x \in X_\sigma$ such that for each $n \geq 1$ there are $a \in \mathcal{A}_i$ and $k \geq 1$ with $x[-n, n] \prec \sigma^k(a)$. Observe that each set Y_i is S-invariant. It is straightforward to check that

$$X_\sigma = \bigsqcup_{i=1}^{m} Y_i. \tag{6.2}$$

Since the matrices F_i are primitive and $\sigma(\mathcal{A}_i) \subset \mathcal{A}_i^+$, we get that for each $i = 1, \ldots, s$, the set Y_i is closed and the system (Y_i, S) is minimal.

To study second order ergodic theorems, we need to single out an infinite invariant measure. In view of (6.2), each invariant ergodic measure μ is supported on one of the sets $Y_1, \ldots, Y_m$. However, not every set Y_i, $i = s+1, \ldots, m$, supports a *"natural"* infinite invariant measure. We do not state the complete description of such invariant measures here, but refer the reader to [BKMS1].

Our approach is based on Theorem 5.4, which we apply to the self-similar substitution system on the line $\mathbb{R}$, arising from the symbolic substitution system (X_σ, S) as a suspension flow. For this to exist, however, it is necessary and sufficient that the substitution matrix M_σ should have a strictly positive left eigenvector, whose components will serve as the lengths of the prototiles. This is a significant restriction: it is known that a strictly positive left eigenvector for the matrix M_σ in the form (6.1) exists if and only if all the matrices $F_1, \ldots, F_s$, corresponding to the minimal components, have the same spectral radius, which is strictly greater than the spectral radii of all the remaining diagonal blocks $F_{s+1}, \ldots, F_m$, see [G, Th.III.6, p.92].

Standing assumption. We will assume for simplicity that

$$M_\sigma = \begin{pmatrix} A & C \\ 0 & B \end{pmatrix}, \tag{6.3}$$

where A and B are primitive, with $\rho(A) > \rho(B) > 1$, and C is non-zero. We expect that our method works in the more general case, when M_σ has a strictly positive left eigenvector, but we have not verified the details.

Under our standing assumption, plus a technical condition stated below, the system (X_σ, S) has a unique, up to scaling, invariant measure that is positive and finite on at least one open set, and this measure is infinite σ-finite. This follows from Corollary 5.6 in [BKMS1] in the case when the substitution system is non-periodic. The non-periodicity was needed to ensure the recognizability property [BKM, Theorem 5.17], however, it is possible to extend the proof of Theorem 5.17 from [BKM] to the needed generality. Here, instead, we use a more direct approach, relating the substitution system to the tiling system.

First we need a technical lemma. Given two sequences $\{x_n\}$ and $\{y_n\}$ of reals, the notation $x_n \approx y_n$ means that $x_n/y_n \to 1$ as $n \to \infty$.

Lemma 6.2. *Let $\mathcal{A} = \{1, 2, \dots, N\}$ be a finite alphabet and $\sigma : \mathcal{A} \to \mathcal{A}^+$ a substitution with the substitution matrix of the form (6.3). Assume that the matrices A and B are primitive and $\rho(A) > \rho(B) > 1$. Then*

$$|\sigma^k(i)| \approx \xi_i \rho(A)^k, \quad i = 1, \dots, N, \tag{6.4}$$

where $\overline{\xi} = (\xi_i)_{i=1}^N$ is a left Perron-Frobenius eigenvector for $M = M_\sigma$, i.e.

$$[\xi_1 \dots \xi_N] M = \rho(A)[\xi_1 \dots \xi_N].$$

Proof. We have $|\sigma^k(i)| = \langle M^k \mathbf{e}_i, \overline{1} \rangle$ where $\mathbf{e}_i$ is the i-th unit vector and $\overline{1} = [1 \dots 1]^t$. Asymptotics of the entries of powers of a non-negative (not necessarily irreducible) matrix are known. It follows e.g. from Theorem (9.4) in [S] that there exist $\xi_i > 0$ such that $|\sigma^k(i)| \approx \xi_i \rho(A)^k$, $i \le N$. It remains to show that $\overline{\xi} = \{\xi_i\}$ is a left eigenvector.

Notice that

$$M^{k+1}\mathbf{e}_j = \sum_{i=1}^N M(i,j) M^k \mathbf{e}_i.$$

Hence

$$\langle M^{k+1}\mathbf{e}_j, \overline{1} \rangle = \sum_{i=1}^N \langle M^k \mathbf{e}_i, \overline{1} \rangle M(i,j),$$

which implies

$$\xi_j \rho(A)^{k+1} \approx \sum_{i=1}^N \xi_i \rho(A)^k M(i,j) \text{ as } k \to \infty.$$

This implies that $\overline{\xi}$ is a left eigenvector for M, as desired. □

Definition 6.3. Set $\lambda = \rho(A)$. Let $\overline{\xi}$ be the left eigenvector for the matrix M_σ in the lemma above, satisfying (6.4). For each letter $a \in \mathcal{A}$, denote by I_a the interval of length ξ_a centered at the origin. We will consider these intervals as tiles in $\mathbb{R}$, labeled by their letters. Set $\varphi(x) = \lambda x$. Define the tile substitution $\mathcal{G}$ on the tiles $\{I_a\}_{a \in \mathcal{A}}$ as follows. Consider the inflated tile $\varphi(I_v) = \lambda I_v$. Since

$$\text{length}(\lambda I_v) = \lambda \xi_v = \sum_{w \in \mathcal{A}} M(w,v) \xi_w,$$

we can subdivide the interval λI_v into the intervals $\{I_w\}$ according to the sequence of all the letters of $\sigma(v)$. Define $\mathcal{G}(I_v)$ as the collection of the corresponding translates of the intervals $\{I_w\}_{w \in \sigma(v)}$. We will call $\mathcal{G}$ the *tile substitution associated to* σ and denote by $\Omega_{\mathcal{G}}$ the corresponding tiling space.

Denote by $\mathcal{B}$ the set of letters corresponding to the matrix B.

Lemma 6.4. *(i) The tiling substitution $\mathbb{R}$-action $(\Omega_{\mathcal{G}}, \mathbb{R})$ is isomorphic (canonically topologically conjugate) to the suspension flow over the symbolic substitution $\mathbb{Z}$-action (X_σ, S), with the "roof function" equal to the constant $\xi_j > 0$ on the cylinder sets $[j]$ for $j \in \mathcal{A}$.*

(ii) Assume that the substitution $\mathcal{G}$ satisfies the conditions of Section 2.3. Then there is a unique infinite (σ-finite) invariant measure ν for the system (X_σ, S), normalized so that

$$\sum_{b\in\mathcal{B}} \xi_b \nu([b]) = 1.$$

This measure may be identified with the transverse measure $\mu^{\rm tr}$ of the invariant measure μ for the system $(\Omega_{\mathcal{G}}, \mathbb{R})$, normalized so that $\mu(\Omega_{\mathcal{B}}) = 1$, where $\Omega_{\mathcal{B}}$ is the set of tilings from $\Omega_{\mathcal{G}}$ having a tile of type $\mathcal{B}$ covering the origin.

Proof. This follows from definitions and the results of [CS]. We just observe that the transversal of $\Omega_{\mathcal{G}}$ may be naturally identified with X_σ, and transversal measures correspond to invariant measures for (X_σ, S). □

For the technical assumptions from Section 2.3 to hold, it is enough that

$$\forall\, b \in \mathcal{B},\ \sigma(b) \text{ starts and ends with a letter from } \mathcal{B}, \tag{6.5}$$

and

$$\forall\, b \in \mathcal{B},\ \exists\, k \in \mathbb{N} \text{ such that } \sigma^k(b) \text{ has at least one "interior" letter from } \mathcal{B}. \tag{6.6}$$

Definition 6.5. We will call a substitution σ *admissible* if it has the form (6.3), with $\rho(A) > \rho(B) > 1$, and both (6.5) and (6.6) are satisfied.

Remark 6.6. Actually, condition (6.5) may be omitted: it implies the "non-periodic border condition", see Definition 2.8, which was needed for recognizability of non-periodic tilings. In fact, in the setting of one-dimensional self-similar tiling substutions, the proof of recognizability from [CS] works without it.

Let (X, T, ν) be an infinite ergodic measure-preserving transformation. The system is called *conservative* if it has no wandering sets of positive measure, i.e. there is no set $W \subset X$ with $\nu(W) > 0$ and $W \cap T^{-n}W = \emptyset$ for every $n \geq 1$. We need conservativity of our systems, since we will consider one-sided averages for the substitution $\mathbb{Z}$-action.

Lemma 6.7. *The substitution dynamical system with an infinite invariant measure (X_σ, ν, S) , corresponding to an admissible substitution σ, is conservative.*

Proof. We will use Maharam's recurrence theorem (see [A, 1.1.7]), which says that if there exists a subset Y of finite measure, such that $X_\sigma = \bigcup_{n=0}^{\infty} S^{-n}Y \bmod \nu$, then S is conservative. Let Y be the set of sequences

$(y_n)_{n\in\mathbb{Z}} \in X_\sigma$ such that $y_1 \in \mathcal{B}$. Then $\nu(Y) < \infty$ by Lemma 6.4. We have $y \notin \bigcup_{n=0}^{\infty} S^{-n}Y$ if and only if there exists $k \in \mathbb{Z}$ such that $y_n \in \mathcal{A}\setminus\mathcal{B}$ for all $n > k$. Since ν is supported on the set of sequences which contain at least one $\mathcal{B}$-symbol, it suffices to show that

$$\nu(Y_0) = 0, \quad \text{where } \; Y_0 = \{x \in X_\sigma : \; x_n \in \mathcal{A}\setminus\mathcal{B} \text{ for all } n > 0 \text{ and } x_0 \in \mathcal{B}\}.$$

For every $y \in Y_0$ and $n > 0$ there exist $b \in \mathcal{B}$ and $i = 0, \dots, |\sigma^n(b)| - 1$ such that $y \in S^i[\sigma^n(b)]$, where $[\sigma^n(b)] = \{x \in X_\sigma : x[0, |\sigma^n(b)| - 1] = \sigma^n(b)\}$. Since every word $\sigma^n(b)$, $b \in \mathcal{B}$, ends with a letter from $\mathcal{B}$, we immediately get that $i = |\sigma^n(b)| - 1$. It follows that $Y_0 \subset \bigcup_{b\in\mathcal{B}} \bigcap_{n\ge 1} S^{|\sigma^n(b)|-1}[\sigma^n(b)]$.

Since the measure ν is non-atomic, we have that

$$\nu(S^{|\sigma^n(b)|-1}[\sigma^n(b)]) = \nu([\sigma^n(b)]) \to 0 \text{ as } n \to \infty.$$

This yields the result. □

The following simple “folklore” lemma gives the so-called “accordion” representation of words from X_σ.

Lemma 6.8. *Let $x \in X_\sigma$ and $n \ge 1$. Then*

$$x[1,n] = u_0\sigma(u_1)\sigma^2(u_2)\dots\sigma^m(u_m)\sigma^m(v_m)\sigma^{m-1}(v_{m-1})\dots\sigma(v_1)v_0, \tag{6.7}$$

where $m \ge 1$ and u_i, v_j, $i,j = 1,\dots,m$, are subwords (possibly empty) of $\sigma(a)$, $a \in \mathcal{A}$. However, at least one of u_m, v_m is nonempty.

Proof. Set $w = x[1,n]$. By the definition of X_σ, we can choose $a \in \mathcal{A}$ and the minimal $k \in \mathcal{N}$ such that $w \prec \sigma^k(a)$. Writing $\sigma^{k-1}(a) = a_1 \dots a_m$ we obtain that $w \prec \sigma(a_1)\dots\sigma(a_m)$, hence

$$w = u_0\sigma(w^{(1)})v_0,$$

where $w^{(1)}$ is a subword $\sigma^{k-1}(a)$ (possibly empty), u_0 is a suffix of some $\sigma(a_i)$ (possibly empty), and v_0 is a prefix of some $\sigma(a_j)$ (possibly empty). Repeating this process with $w^{(1)}$, etc. by induction, we obtain the desired representation (6.7). □

For a word $w \in \mathcal{A}^+$ we define its “population vector” by $\overline{\ell}(w) = (\ell_i(w))_{i=1}^N$ where $\ell_i(w)$ is the number of symbols i in the word w. For $w \in \mathcal{A}^+$ denote

$$|w|_\mathcal{T} := \langle \overline{\ell}(w), \overline{\xi} \rangle \tag{6.8}$$

and call this quantity the *tiling length* of the word w. Note that $\overline{\ell}(\sigma(w)) = M_\sigma\overline{\ell}(w)$ by definition of the substitution matrix.

Lemma 6.9. *Let σ be an admissible substitution. Then for any $x \in X_\sigma$ we have*

$$\lim_{n\to\infty} \frac{|x[1,n]|_\mathcal{T}}{n} = 1.$$

Proof. Given $x \in X_\sigma$ and $n \geq 1$, consider the accordion representation (6.7). Note that for all i,

$$|u_i|, |v_i| \leq \max_a |\sigma(a)| =: L_{\max}.$$

Recall that for each letter $a \in \mathcal{A}$ we have $|\sigma^k(a)| \approx \xi_a \lambda^k = \lambda^k |a|_{\mathcal{T}}$ by Lemma 6.2. Hence

$$|\sigma^k(u)| \approx \lambda^k |u|_{\mathcal{T}}, \quad k \to \infty, \tag{6.9}$$

uniformly for all u with $|u| \leq L_{\max}$. Note also

$$\langle \overline{\ell}(\sigma^k(j)), \overline{\xi} \rangle = \langle M_\sigma^k \mathbf{e}_j, \overline{\xi} \rangle = \langle \mathbf{e}_j, (M_\sigma^t)^k \overline{\xi} \rangle = \langle \mathbf{e}_j, \lambda^k \overline{\xi} \rangle = \lambda^k \xi_j,$$

which yields $|\sigma^k(u)|_{\mathcal{T}} = \lambda^k |u|_{\mathcal{T}}$. Using the accordion representation of $x[1,n]$, we obtain

$$\frac{|x[1,n]|_{\mathcal{T}}}{n} = \frac{\sum_{i=0}^m \lambda^i |u_i|_{\mathcal{T}} + \sum_{i=0}^m \lambda^i |v_i|_{\mathcal{T}}}{\sum_{i=0}^m |\sigma^i(u_i)| + \sum_{i=0}^m |\sigma^i(v_i)|}.$$

Now the desired statement follows from (6.9) and the fact that $m \to \infty$ as $n \to \infty$ and at least one of u_m, v_m is nonempty. □

The next lemma gives an upper bound for the number of $\mathcal{B}$-tiles in the interval $[0,t]$. Recall that $\Omega_{\mathcal{G}}$ is the tiling space of $\mathcal{G}$, the tile substitution associated to σ. Given $\mathcal{T} \in \Omega_{\mathcal{G}}$, denote by $N_{\mathcal{T}}(\mathcal{B},t)$ the total number of $\mathcal{B}$-tiles of $\mathcal{T}$, contained (completely) in the interval $[0,t]$.

Lemma 6.10. *There exists a constant $K > 0$ such that for every $\mathcal{T} \in \Omega_{\mathcal{G}}$ and $t > 0$, we have $N_{\mathcal{T}}(\mathcal{B},t) \leq Kt^\alpha$, where $\alpha = \log(\rho(B))/\log(\rho(A))$.*

Proof. Since the inequality will persist (with a slightly larger constant K) if we shift $\mathcal{T}$ by a fixed vector, we can assume that $\mathcal{T}$ belongs to the transversal of $\Omega_{\mathcal{G}}$. For every integer $s > 0$, find a tiling $\mathcal{T}_s$ such that $\mathcal{G}^s(\mathcal{T}_s) = \mathcal{T}$, see Proposition 2.7. Choose an integer $k > 0$ such that $\rho(A)^k \xi_i > 2$ for every $i \in \mathcal{A}$. Let T_s be the tile of $\mathcal{T}_s$ containing the origin; it is centered at the origin for all s: since $\mathcal{T}$ is in the transversal, all T_s are actual prototiles, see Definition 6.3. Then the interval $[0, \rho(A)^s]$ is covered by the patch $\mathcal{G}^{s+k}(T_s)$ by our choice of k. Thus, $N_{\mathcal{T}}(\mathcal{B}, \rho(A)^s)$ does not exceed the number of occurrences of $\mathcal{B}$-tiles in the patch $\mathcal{G}^{s+k}(T_s)$. By the Perron-Frobenius theorem applied to the primitive matrix B, the number of $\mathcal{B}$-tiles in $\mathcal{G}^{s+k}(T_s)$ asymptotically grows not faster than $K\rho(B)^s$ for some constant K independent of s. The constant K can be adjusted so that $N_{\mathcal{T}}(\mathcal{B}, \rho(A)^s) \leq K\rho(B)^s$ for every tiling $\mathcal{T}$ and every positive real number s. Setting $t = \rho(A)^s$, noting that $\rho(A)^\alpha = \rho(B)$ and adjusting the constant again, we obtain the desired inequality for all $t > 0$. □

Now we are ready to prove the main result on substitutions. It will be convenient to write elements of X_σ as $(x(n))_{n \in \mathbb{Z}}$.

Theorem 6.11. *Let $\sigma : \mathcal{A} \to \mathcal{A}^+$ be an admissible substitution, see Definition 6.5. Let ν be the infinite invariant measure on X_σ from Lemma 6.4.*

Then for every function $f \in L^1(X_\sigma, \nu)$ and ν-a.e. $x \in X_\sigma$, we have that

$$\int_{X_\sigma} f(y) d\nu(y) = \lim_{n\to\infty} \frac{1}{\log(n)} \sum_{k=1}^{n} \frac{\sum_{i=0}^{k-1} f(S^i x)}{ck^\alpha} \frac{1}{k},$$

where $\alpha = \log(\rho(B))/\log(\rho(A))$ and $c > 0$ is the average density of the graph-directed sets associated with $\mathcal{G}$.

Proof. (1) Consider the one-dimensional tile substitution system $(\Omega_\mathcal{G}, \mathbb{R})$ associated with $\mathcal{G}$, see Definition 6.3 and Lemma 6.4. Denote by μ the (unique) translation-invariant measure on $\Omega_\mathcal{G}$ corresponding to ν. Recall that X_σ can be identified with the transversal of $\Omega_\mathcal{G}$. More precisely, for $x \in X_\sigma$, let $\mathcal{T}(x) \in \Omega_\mathcal{G}$ be the tiling, which has the prototile $I_{x(0)}$ as its tile, and the other tiles are $I_{x(n)} + y_n$, so that the left endpoint of $I_{x(n+1)} + y_{n+1}$ is the right endpoint of $I_{x(n)} + y_n$ for all $n \in \mathbb{Z}$.

(2) In view of Theorem 4 in [Fi1], which is a one-sided version of Lemma 5.3, it is enough to establish the result for a single function $f \in L^1(X_\sigma, \nu)$ with $\int f d\nu \neq 0$. Theorem 4 from [Fi1] was established under the assumption that the system is conservative, so we use Lemma 6.7 here. Consider the function f on X_σ given by: $f(x) = 1$ if $x(1) \in \mathcal{B}$ and $f(x) = 0$ otherwise.

Let F be the function on $\Omega_\mathcal{G}$ such that $F(\mathcal{T}) = 1/\xi_i$ if the tile of $\mathcal{T}$ containing the origin is a translate of I_i for some $i \in \mathcal{B}$, and $F(\mathcal{T}) = 0$ otherwise. This is well-defined μ-a.e. Repeating the arguments of Theorem 5.4, we obtain that for μ-a.e. tiling $\mathcal{T} \in \Omega_\mathcal{G}$,

$$\lim_{t\to\infty} \frac{1}{\log(t)} \int_0^t \frac{\int_0^R F(\mathcal{T} - u)\, du}{R^{\alpha+1}} dR = c \int_{\Omega_\mathcal{G}} F(\mathcal{S})\, d\mu(\mathcal{S}) := \theta > 0, \tag{6.10}$$

where $\alpha = \log(\rho(B))/\log(\rho(A))$ and $c > 0$ is the average density of the graph-directed sets associated with $\mathcal{G}$.

Recall that $N_\mathcal{T}(\mathcal{B}, R)$ is the number of $\mathcal{B}$-tiles contained in $[0, R]$. Thus,

$$N_\mathcal{T}(\mathcal{B}, R) \le \int_0^R F(\mathcal{T} - u)\, du \le N_\mathcal{T}(\mathcal{B}, R) + 1.$$

Since μ is (locally) a product of the transverse measure ν and the Lebesgue measure on $\mathbb{R}$, it follows from Equation (6.10) that for ν-a.e. $x \in X_\sigma$,

$$\lim_{t\to\infty} \frac{1}{\log(t)} \int_0^t \frac{N_{\mathcal{T}(x)}(\mathcal{B}, R)}{R^{\alpha+1}} dR = \theta. \tag{6.11}$$

(3) We want to show that

$$\theta = \lim_{n\to\infty} \frac{1}{\log(n)} \sum_{k=1}^{n} \frac{\sum_{i=0}^{k-1} f(S^i x)}{k^{\alpha+1}}$$

for all $x \in X_\sigma$ satisfying Equation (6.11). Denote by $\ell_{\mathcal{B}}(w)$ the number of $\mathcal{B}$-letters in a word $w \in \mathcal{A}^+$, and observe that

$$\sum_{i=0}^{k-1} f(S^i x) = \ell_{\mathcal{B}}(x[1,k]).$$

Note that (6.11) is equivalent to

$$\lim_{n\to\infty} \frac{1}{\log(n)} \sum_{k=1}^{n} \frac{N_{\mathcal{T}(x)}(\mathcal{B},k)}{k^{\alpha+1}} = \theta,$$

so we just need to estimate $|N_{\mathcal{T}(x)}(\mathcal{B},k) - \ell_{\mathcal{B}}(x[1,k])|$. We claim that

$$\ell_{\mathcal{B}}(x[1,k]) = N_{\mathcal{B}}(\mathcal{T}(x), R_k), \quad \text{where} \quad R_k = |x[1,k]|_{\mathcal{T}} + (1/2)\xi_{x(0)}. \tag{6.12}$$

Indeed, the left-hand side represents the number of $\mathcal{B}$-letters in $x[1,k]$ and the right-hand side equals the number of $\mathcal{B}$-tiles, (completely) contained in the interval $[0, R_k]$. According to the definition of $\mathcal{T}(x)$ at the beginning of the proof, it has the prototile $I_{x(0)}$ centered at the origin, so that half of its length $(1/2)\xi_{x(0)}$ is in $[0, R_k]$. After that the sequence of tiles which fits in $[0, R_k]$ exactly corresponds to $x[1,k]$, by the definition of the tile length. Thus, both sides of (6.12) count the same quantity.

Now, by Lemma 6.9 we have $R_k \approx k$, hence $R_k = k + o(k)$, as $k \to \infty$, using the standard $o(\cdot)$ notation. In view of (6.12) and Lemma 6.10, we have

$$\begin{aligned} |N_{\mathcal{T}(x)}(\mathcal{B},k) - \ell_{\mathcal{B}}(x[1,k])| &= |N_{\mathcal{T}(x)}(\mathcal{B},k) - N_{\mathcal{T}(x)}(\mathcal{B},R_k)| \\ &\le K|R_k - k|^\alpha + 1 = o(k^\alpha). \end{aligned}$$

Indeed, $N_{\mathcal{T}(x)}(\mathcal{B},R_k) - N_{\mathcal{T}(x)}(\mathcal{B},k)$ equals the number of $\mathcal{B}$-tiles of $\mathcal{T}(x)$ in the interval $[k, R_k]$ (assume that $k \le R_k$ for definiteness) plus one, if a $\mathcal{B}$-tile contains $k \in \mathbb{R}$ in its interior, and the number of $\mathcal{B}$-tiles of $\mathcal{T}(x)$ in the interval $[k, R_k]$ equals $N_{\mathcal{B}}(\mathcal{T}(x) - k, R_k - k)$, to which we can apply Lemma 6.10. Thus,

$$\begin{aligned} \lim_{n\to\infty} \frac{1}{\log(n)} \sum_{k=1}^{n} \frac{\sum_{i=0}^{k-1} f(S^i x)}{k^{\alpha+1}} &= \lim_{n\to\infty} \frac{1}{\log(n)} \sum_{k=1}^{n} \frac{\ell_{\mathcal{B}}(x[1,k])}{k^{\alpha+1}} \\ &= \lim_{n\to\infty} \frac{1}{\log(n)} \sum_{k=1}^{n} \frac{N_{\mathcal{T}(x)}(\mathcal{B},k)}{k^{\alpha+1}} \\ &+ \lim_{n\to\infty} \frac{1}{\log(n)} \sum_{k=1}^{n} \frac{o(k^\alpha)}{k^{\alpha+1}} = \theta + 0, \end{aligned}$$

as desired. Noticing that

$$\theta = c \int_{\Omega_{\mathcal{G}}} F\, d\mu = c \sum_{b\in\mathcal{B}} \nu([b]) = c \int_{X_\sigma} f\, d\nu$$

by Lemma 6.4, we get the result. □

Remark 6.12. We note that the parameters α and c appearing in the second order ergodic theorem are invariants of the measure-theoretical isomorphism between infinite measure preserving systems. This is immediate, since there exists $f \in L^1(X_\sigma, \nu)$ such that $\int_{X_\sigma} f d\nu$ is positive and finite.

As an example, consider two symbolic substitution systems on the alphabet $\mathcal{A} = \{0, 1\}$ given by $\sigma_1(0) = 0^3$ (three zeros), $\sigma_1(1) = 101$; and $\sigma_2(0) = 0^9$, $\sigma_2(1) = 1^4 0 1^4$. Then, $\alpha_1 = \log(2)/\log(3)$, whereas $\alpha_2 = \log(8)/\log(9)$. Since $\alpha_1 \neq \alpha_2$, these systems cannot be measure-theoretically isomorphic with respect to the invariant infinite measures, and hence, cannot be topologically conjugate.

The parameter c can also be used to distinguish substitution systems, although the computation is more involved. For example, consider for $k = 0, \ldots, 3$ the substitutions $\sigma_k(0) = 0^9$ and $\sigma_k(1) = 10^k 10^{6-k} 1$. For all of them we have $\alpha = 1/2$, but the average densities of the corresponding graph-directed sets are likely to be different, which would imply that the substitution dynamical systems associated with σ_k are pairwise non-isomorphic.

As a corollary of Theorem 6.11, we establish that almost every sequence in X_σ admits an "α-dimensional density".

Corollary 6.13. *Let (X_σ, ν, S) be a substitution system satisfying the assumptions of Theorem 6.11. Then for every letter $b \in \mathcal{B}$, the limit*

$$\lim_{n\to\infty} \frac{1}{\log(n)} \sum_{1\le k\le n,\ x_k=b} \frac{1}{k^\alpha}$$

exists and equals to $\alpha \cdot c \cdot \nu([b])$ for ν-a.e. $x = (x_k) \in X_\sigma$.

Proof. We will use the same notation as in the proof of Theorem 6.11. Fix a letter $b \in \mathcal{B}$. Consider the function $f : X_\sigma \to \mathbb{R}$ such that $f(x) = 1$ if $x_0 = b$ and $f(x) = 0$ otherwise. Given a sequence $x \in X_\sigma$, denote by $\ell_b(x, k) = \ell_b(x[1, k])$ the number of occurrences of the symbol b in the word $x[1, k]$. Theorem 6.11 implies that

$$\lim_{n\to\infty} \frac{1}{\log(n)} \sum_{k=1}^{\infty} \frac{\ell_b(x, k)}{k^{\alpha+1}} = c\nu([b]) \tag{6.13}$$

for ν-a.e. $x \in X_\sigma$. Fix a sequence $x \in X_\sigma$ satisfying Equation (6.13). Using summation by parts, we get

$$\sum_{k=1}^{n} \frac{f(S^k x)}{k^\alpha} = \sum_{k=1}^{n-1} \ell_b(x, k) \left(\frac{1}{k^\alpha} - \frac{1}{(k+1)^\alpha} \right) + \frac{\ell_b(x, n)}{n^\alpha}.$$

Lemmas 6.9 and 6.10 imply that $\ell_b(x,n)/\alpha^d$ is uniformly bounded in n. Notice that $\left(\frac{1}{k^\alpha} - \frac{1}{(k+1)^\alpha}\right) \approx \frac{\alpha}{k^{\alpha+1}}$. Thus, (6.13) yields

$$\begin{aligned} \lim_{n\to\infty} \frac{1}{\log(n)} \sum_{k=1}^{n} \frac{f(S^k x)}{k^\alpha} &= \lim_{n\to\infty} \frac{1}{\log(n)} \sum_{k=1}^{n} \ell_b(x,k) \frac{d}{k^{\alpha+1}} \\ &= \alpha \int f d\nu = \alpha \cdot c \cdot \nu([b]) \end{aligned}$$

for ν-a.e. $x \in X_\sigma$. □

This may be compared with a result of [Be], which implies that all "morphic" sequences x have the *logarithmic frequency* of letters. This means that for $a \in \mathcal{A}$ the following limit exists:

$$\lim_{n\to\infty} \frac{1}{\log(n)} \sum_{k=1,\, x_k=a}^{n} \frac{1}{k}\,.$$

For our substitutions it is immediate that the logarithmic frequency of $b \in \mathcal{B}$ is zero for all $x \in X_\sigma$, since already the ordinary frequency $\lim_{n\to\infty} \frac{1}{n}\#\{k \le n :\ x_k = b\}$ equals zero.

7. Examples and open questions

In this section we consider a few examples of tiling and substitution systems and determine the parameter d appearing in the second order ergodic theorem.

Example 7.1. ("Cantor" substitution). Let $\mathcal{A} = \{0,1\}$ and $\sigma(0) = 000$, $\sigma(1) = 101$. Consider the substitution system (X_σ, S) associated to σ. Observe that the graph-directed set of σ is the middle-third Cantor set. The dynamical system (X_σ, S) admits a unique ergodic measure μ on X_σ with the property $\mu([1]) = 1$. Then Theorem 6.11 holds for the system (X_σ, S) with parameters $\alpha = \log(2)/\log(3)$ and $c > 0$, where c is the average density of the middle-third Cantor set. We refer the reader to Theorem 6.6 of [Fa] for the numerical value of c. We mention that the second order theorem for the system (X_σ, S) was originally established (it was actually the main result) by A. Fisher [Fi1].

Example 7.2. Ergodic measures of the following dynamical system were described in [CS]. Suppose that the alphabet $\mathcal{A}$ consists of two 1×1 squares on the plane labeled by 0 and 1. Denote the tiling substitution by

$$\boxed{0} \mapsto \begin{array}{|c|c|c|} \hline 0 & 0 & 0 \\ \hline 0 & 0 & 0 \\ \hline 0 & 0 & 0 \\ \hline \end{array} \qquad \text{and} \qquad \boxed{1} \mapsto \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 1 & 0 & 1 \\ \hline 1 & 1 & 1 \\ \hline \end{array}$$

The associated graph-directed set is the Sierpiński carpet. So Theorem 5.4 with parameter $\alpha = \log(8)/\log(3)$ applies to this tiling system.

Example 7.3. In this example, we consider a tiling dynamical systems with prototiles having fractal boundaries. Our example is a modification of the system described in [So1, Section 7.2] and belonging to the family of tilings constructed in [K, Section 6].

Let $r \approx .34115 + 1.1616i$ be a root of the equation $x^3 + x + 1 = 0$. Let T_a, T_b, and T_c be sets (prototiles) as described in Lemma 7.7 of [So1]. We use the same notation as in [So1].

We note that T_a, T_b, and T_c are compact subsets of $\mathbb{C}$. Set $\theta(z) = rz$. Then $\theta(T_a) = T_b$; $\theta(T_b)$ is the union of a translation of T_b and of T_c; and $\theta(T_c)$ is a translation of T_a. This subdivision rule uniquely determines a tile substitution Θ.

Set $\mathcal{G} = \Theta^2$. Thus, $\mathcal{G}(T_a)$ is the union of a translation of T_b and of T_c; $\mathcal{G}(T_b)$ is the union of a translation of T_b, of T_c, and of T_a; and $\mathcal{G}(T_c)$ is a copy of T_b. Set $\varphi(z) = r^2 z$. Then after the "realification" of $\mathbb{C}$, the map φ can be represented as $\varphi((x,y)^T) = \lambda \cdot O \cdot (x,y)^T$, where $\lambda = |r^2|$ and

$$O = \begin{pmatrix} \beta & -\gamma \\ \gamma & \beta \end{pmatrix}$$

with $\beta + \gamma i = r^2/|r^2|$. Then λ is the expansion constant of φ.

Assuming that the tiles T_a, T_b, and T_c are colored in white color, denote by S_a, S_b, and S_c their respective copies colored in a dark color. Extend the tile substitution $\mathcal{G}$ on $\{S_a, S_b, S_c\}$ as follows: S_a is mapped into a union of S_b and S_c in the same way as T_a; S_c is mapped into a copy of S_b as T_c; and the tile S_b is mapped into a union of S_a, S_b, and S_c exactly as T_b, but with the tile S_b being replaced by the tile T_b (of the same shape but of a different color).

Denote by $\mathcal{A}$ the set of prototiles $\{T_a, T_b, T_c, S_a, S_b, S_c\}$. Consider the tiling dynamical system $(\Omega_{\mathcal{G}}, \mathbb{R}^2)$ associated to $\mathcal{G} : \mathcal{A} \to \mathcal{A}^+$. We notice that this system has a unique minimal component determined by tiles of white color, see [CS, Lemma 2.10]. Observe that the minimal component is non-periodic, see [So1, Section 7.2]. Hence, the substitution $\mathcal{G}$ satisfies all the conditions of Theorem 2.12 yielding that this system admits a unique "natural" infinite invariant measure μ up to scaling.

The substitution matrix $M(\mathcal{G})$ is given by

$$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 \end{pmatrix}$$

It follows that the matrix B is equal to

$$\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}.$$

Since $\rho(B) \approx 1.618$ and the expansion constant of $\mathcal{G}$ is $\lambda \approx 1.466$, we get that $\alpha = \log(\rho(B))/\log(\lambda) > 1$. Hence, the second order theorem (Theorem 5.4) with parameter $\alpha \approx 1.258$ applies to the system $(\Omega_{\mathcal{G}}, \mathbb{R}^2)$.

Example 7.4. This example is a non-minimal extension of the well-known Rauzy tiling [R].

We start with the Rauzy tiling itself. Let $r \approx -0.7771845 + 1.11514i$ be the complex root of the equation $1 - r - r^2 - r^3 = 0$. The tiles may be described using digit expansions in the base of r. There are three prototiles T_a, T_b, and T_c, which may be represented as follows:

$$T := \Big\{ \sum_{n=0}^{\infty} a_n r^{-n} : \ a_n \in \{0,1\},\ a_n a_{n+1} a_{n+2} \neq 111 \text{ for all } n \Big\}.$$

Then

$$T_a := r^{-1}T,\ T_b := 1 + r^{-2}T,\ T_c := 1 + r^{-1} + r^{-3}T.$$

Clearly, $rT_a = T_a \cup T_b \cup T_c$, $rT_b = r + T_a$, and $rT_c = r + T_b$. This determines the substitution rule. (Strictly speaking, these prototiles do not satisfy our definition, since T_b and T_c do not contain the origin in the interior of their support, but this can be easily rectified, translating the tiles. However, the given form of the tiles is more convenient.) All the tiles of the Rauzy tiling can also be described using base r expansions: for any *finite* sum $z = \sum_{n=-N}^{-1} a_n r^{-n}$ with the property that $a_n \in \{0,1\}$, $a_n a_{n+1} a_{n+2} \neq 111$ for all n, we have $z + T_a \in \mathcal{T}$ in all cases, $z + T_b \in \mathcal{T}$ iff $a_{-2}a_{-1} \neq 11$, and $z + T_c \in \mathcal{T}$ iff $a_{-1} \neq 1$.

Now consider the "extended" tiling system, with the prototiles T_a, T_b, T_c and S_a, S_b, which have the same support as T_a, T_b respectively, but have a different color (label). The substitution acts as before on T_a, T_b, T_c, and

$$rS_a = S_a \cup S_b \cup T_c, \quad rS_b = 1 + S_a.$$

The matrix of the substitution is $M(\mathcal{G}) = \begin{pmatrix} A & C \\ 0 & B \end{pmatrix}$, where

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \quad C = \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

The expansion $\lambda = |r| \approx 1.3562$ is the same as for the Rauzy tiling, and $\rho(B) \approx 1.618$ is the golden ratio. All the assumptions from Section SectionAssumptions are easily verified. We get $\alpha = \log(\rho(B))/\log(\lambda) \approx 1.57935 > 1$, so Theorem 5.4 applies. The figure shows the "cantorization" of the tiling, so it gives an idea of both "large-scale" structure of the tiling and the "small-scale" structure of the graph-directed sets.

It is interesting to note that the "cantorization" of the tiling has a simple description using base r expansions: instead of all expansions using the digits (a_n) with 111 forbidden, one should consider all expansion with the sequence of digits from the "golden mean" shift, that is, 11 is forbidden.



Figure 1. "Cantorization" of the tiling

7.1. **Open questions.** 1. We had to impose some technical conditions on the substitution to prove the second order ergodic theorem. For example, we do not know if it holds for the following substitution on $\{0,1,2\}$:

$$0 \mapsto 00000, \quad 1 \mapsto 1111, \quad 2 \mapsto 20212.$$

The matrix of the substitution is $M_\sigma = \begin{pmatrix} 5 & 0 & 1 \\ 0 & 4 & 1 \\ 0 & 0 & 3 \end{pmatrix}$. Thus, there is an infinite (σ-finite) invariant measure positive and finite on cylinder sets containing 2, however, there is no positive left eigenvector, so our methods do not work.

2. We proved that (in appropriate contexts) converge the logarithmic averages of the expressions

$$R^{-d}\int_{B_R} f(\mathcal{T}-u)\,du \quad\text{and}\quad k^{-d}\sum_{i=0}^{k-1} f(S^i x).$$

But one can also view them as random variables (with $\mathcal{T}$ or x taken randomly from the substitution space, according to the invariant measure normalized on the appropriate cylinder set), and inquire whether they converge in distribution as R (resp. k) tend to infinity along a subsequence? For instance, it appears that for the "integer Cantor" substitution from Example 1, we get that $2^{-n}\sum_{i=0}^{3^n-1} f(S^i x)$ tends to the uniform distribution on $[0,1]$ as $n\to\infty$, for f the characteristic function of [1] (and then for all $f\in L^1(X_\sigma,\mu)$ with $\int f\,d\mu=1$).

Acknowledgment. We are grateful to Karl Petersen, who asked whether the second-order ergodic theorem holds for the Sierpiński gasket tiling system and who told us about A. Fisher's paper [Fi1].

References

[A] J. Aaronson, *An introduction to infinite ergodic theory.* Mathematical Surveys and Monographs, 50. American Mathematical Society, Providence, RI, 1997. xii + 284 pp.

[ADF] J. Aaronson, M. Denker, and A. Fisher, *Second order ergodic theorems for ergodic transformations of infinite measure spaces.* Proc. Amer. Math. Soc. 114 (1992), no. 1, 115 – 127.

[AP] J. Anderson and I. Putnam, *Topological invariants for substitution tilings and their associated C^*-algebras.* Ergodic Theory Dynam. Systems **18** (1998), no. 3, 509 – 537.

[Beck] M. Becker, *A ratio ergodic theorem for groups of measure-preserving transformations.* Illinois J. Math. **27** (1983), 562 – 570.

[Be] J. Bell, *Logarithmic frequency in morphic sequences.* J. Theor. Nombres Bordeaux 20 (2008), no. 2, 227 – 241.

[BKM] S. Bezuglyi, J. Kwiatkowski, K. Medynets, *Aperiodic substitution systems and their Bratteli diagrams.* Ergodic Theory Dynam. Systems **29** (2009), no. 1, 37 – 72.

[BKMS1] S. Bezuglyi, J. Kwiatkowski, K. Medynets, B. Solomyak, *Invariant measures on stationary Bratteli diagrams.* Ergodic Theory Dynam. Systems **30** (2010), no. 4, 973 – 1007.

[CS] M.I. Cortez and B. Solomyak, *Invariant measures for non-primitive tiling substitutions.* J. Anal. Math. **115** (2011), 293 – 342.

[E] G. Edgar, *Integral, Probability, and Fractal Measures.* Springer, 1997. x+286pp.

[Fa] K. Falconer, *Techniques in fractal geometry.* John Wiley & Sons, Ltd., Chichester, 1997. xviii+256 pp.

[Fi1] A. Fisher, *Integer Cantor sets and an order-two ergodic theorem.* Ergodic Theory Dynam. Systems **13** (1993), no. 1, 45 – 64.

[G] F. R. Gantmacher, *Applications of the theory of martrices.* Interscience Publishers, Inc., New York, 1959. x+317pp.

[HY] M. Hama and H. Yuasa, *Invariant measures for subshifts arising from substitutions of some primitive components.* Hokkaido Math. J. **40** (2011), no. 2, 279 - 312.

[H] M. Hochman, *A ratio ergodic theorem for multiparameter non-singular actions.* J. Eur. Math. Soc. **12** (2010), no. 2, 365 – 383.

[HuW] W. Hurewicz, and H. Wallman, *Dimension Theory.* Princeton Mathematical Series, v. 4. Princeton University Press, Princeton, N. J., 1941. vii+165 pp.

[K] R. Kenyon, *The construction of self-similar tilings.* Geom. Funct. Anal. **6** (1996), no. 3, 471 – 488.

[LS] F. Ledrappier and O. Sarig, *Fluctuations of ergodic sums for horocycle flows on* $\mathbb{Z}^d$*-covers of finite volume surfaces.* Discrete Contin. Dyn. Syst. **22** (2008), no. 1-2, 247 – 325.

[MW] R.D. Mauldin, S.C. Williams, *Hausdorff dimension in graph directed constructions.* Trans. Amer. Math. Soc. **309** (1988), no. 2, 811 – 829.

[P] B. Praggastis, *Numeration systems and Markov partitions from self similar tilings.* Trans. Amer. Math. Soc. **351** (1999), no. 8, 3315 – 3349.

[RW] C. Radin, M. Wolff, *Space tilings and local isomorphism.* Geom. Dedicata **42** (1992), no. 3, 355 – 360.

[R] G. Rauzy, *Nombres algébriques et substitutions.* Bull. Soc. Math. France **e**110(2) (1982), 147 – 178.

[S] H. Schneider, *The influence of the marked reduced graph of a nonnegative matrix on the Jordan form and on related properties: a survey.* Proceedings of the symposium on operator theory (Athens, 1985). Linear Algebra Appl. **84** (1986), 161 – 189.

[So1] B. Solomyak, *Dynamics of self-similar tilings.* Ergodic Theory Dynam. Systems **17** (1997), no. 3, 695 – 738.

[So2] B. Solomyak, *Nonperiodicity implies unique composition for self-similar translationally finite tilings.* Discrete Comput. Geom. **20** (1998), 265 – 279.

[Y] H. Yuasa, *Invariant measures for the subshifts arising from non-primitive substitutions.* J. D'Analyse Math. **102** (2007), 143 – 180.

Konstantin Medynets, Department of Mathematics, U.S. Naval Academy, Annapolis, MA 21402, USA
E-mail address: `medynets@usna.edu`

Boris Solomyak, Department of Mathematics, University of Washington, Seattle, WA 98195-4350, USA
E-mail address: `solomyak@uw.edu`